

Supplementary Information to the manuscript

Huaier n-butanol Extract Suppresses Proliferation and Metastasis of Gastric Cancer via c-Myc-Bmi1

Axis

Yiping Wang¹, Hang Lv¹, Zhiyuan Xu^{1,2}, Jiancheng Sun², Yixiu Ni², Zhe Chen¹, Xiangdong Cheng^{1,2*}

¹ Key Laboratory of Integrated Traditional Chinese and Western Medicine for Diagnosis and Treatment of Digestive System Tumor, the First Affiliated Hospital of Zhejiang Chinese Medical University, 54 Youdian Road, Hangzhou, 310006, China

² Department of Gastrointestinal Surgery, the First Affiliated Hospital of Zhejiang Chinese Medical University, Hangzhou, Zhejiang, China

To whom correspondence should be addressed: Xiangdong Cheng, Key Laboratory of Integrated Traditional Chinese and Western Medicine for Diagnosis and Treatment of Digestive System Tumor, the First Affiliated Hospital of Zhejiang Chinese Medical University, 54 Youdian Road, Hangzhou, 310006, China. Tel.: Tel: +86-571-87070965; Fax: +86-571-87070965; E-mail: chengxd@zcmu.edu.cn

This file contains supplementary figure legends and two supplementary figures.

Supplementary figure legends

Supplementary figure 1 Full-length Western blots appeared in the main manuscript

The full-length Western blots of Figure 3C and Figure 5A are included.

The rectangle boxes indicate the areas of the Western blots shown in the main figures. MW kDa- molecular weights in kiloDaltons of the marker proteins. AB indicates the antibody used for the blot.

Supplementary figure 2 Full-length Western blots appeared in the main manuscript

The full-length Western blots of Figure 4E are included. The rectangle boxes indicate the areas of the Western blots shown in the main figures. MW kDa: molecular weights in kiloDaltons of the marker proteins. AB indicates the antibody used for the blot.

Supplementary figure 1

Figure 3C





Figure 5A



Supplementary figure 2

Figure 4E

