

Supplemental Table 1. Clinical characteristics of patients with motor functional neurological disorders.

| FND Subject | Age | Gender | Educ. (years) | FND Subtype | Symptom Description | Current Psychiatric Diagnoses | Past Psychiatric Diagnoses | Psychotropic Medications |
|-------------|-----|--------|---------------|-----------------|---|---|--------------------------------|--------------------------|
| 1 | 60 | M | 18 | PNES, FW | Episodes of behavioral arrest with altered consciousness, left arm and leg weakness | Anxiety NOS | MDD, LSD and Opiate Abuse | GBP, MIR |
| 2 | 55 | F | 12 | FMD, FW* | Right arm tremor, right leg weakness | MDD, PD_Ag+ | MDD, Alcohol and Cocaine Abuse | SERT, TZD |
| 3 | 55 | F | 16 | FMD* | Abnormal gait | GAD, Undiff Somatoform | - | ZLP, ECP, GBP |
| 4 | 49 | M | 14 | FW | Left arm and leg weakness | Depression NOS**, Anxiety NOS** | - | - |
| 5 | 36 | F | 12 | FMD, FW | Right arm tremor, left facial weakness | Dysthymia, PD_Ag-, GAD, Undiff Somatoform | MDD | DLX, TZD |
| 6 | 36 | M | 16 | FMD* | Head and bilateral limb tremor | - | - | - |
| 7 | 26 | F | 15 | PNES | Episodes of behavioral arrest with altered consciousness | BPAD II, PD_Ag+, OCD, PTSD | - | GBP, CLP, LRZ, NRT |
| 8 | 42 | F | 18 | FW* | Right hand weakness | MDD, PD_Ag-, GAD, Somatoform Pain | Opiate Abuse | CLP, DVX |
| 9 | 44 | F | 18 | FMD | Abnormal gait | - | Eating Disorder NOS | DLX |
| 10 | 31 | F | 15 | FW [†] | Right arm and leg weakness | PTSD, OCD, Ag | MDD | - |

| | | | | | | | | |
|----|----|---|----|---------------|--|---|----------------------------------|-------------------------------|
| 11 | 50 | F | 14 | FMD, FW*† | Abnormal gait, head and bilateral arm tremors, right leg weakness | PD | - | DLX, CLP, GBP, BSP |
| 12 | 51 | F | 14 | FMD* | Head and four limb tremor, neck dystonia | - | MDD | - |
| 13 | 49 | F | 14 | FMD, FW* | Left facial weakness, head tremor, abnormal gait | MDD, Undiff Somatoform | PTSD | DOX |
| 14 | 26 | F | 16 | FMD, PNES* | Head and bilateral arm tremor, abnormal gait, motor seizures | MDD, PD_Ag+, GAD, Undiff Somatoform | - | CLP, VEN |
| 15 | 30 | F | 14 | FW† | Four limb weakness | MDD, PD_Ag+, PTSD, GAD, Somatization Disorder | MDD | APZ, CLP, PGN, FLX, DZP |
| 16 | 23 | F | 18 | PNES, FW† | Right arm weakness, atonic seizures | Depression NOS, Social Phobia, Alcohol Abuse | - | - |
| 17 | 22 | M | 14 | FW† | Left arm and leg weakness | Depression NOS, PD_Ag- | - | - |
| 18 | 28 | M | 14 | PNES | Episodes of behavioral arrest with altered consciousness, Motor seizures | Ag | Depression NOS, Social Phobia | CLP |
| 19 | 20 | F | 14 | PNES | Motor seizures | PD_Ag+, Social Phobia, Specific Phobia, GAD | Depression NOS | SERT |
| 20 | 50 | F | 12 | PNES | Motor seizures | Depression NOS, | Depression NOS, | ATP, CLP, |

| | | | | | | | | |
|----|----|---|----|-------------------|---|---|--|-------------------------------------|
| | | | | | | PD_Ag-, PTSD | Eating Disorder NOS, Alcohol and Cocaine Abuse | VPA |
| 21 | 27 | F | 18 | PNES | Motor seizures | PD_Ag+ | MDD | CLN |
| 22 | 45 | M | 14 | FMD* | Head and bilateral leg tremor, abnormal gait | MDD | PD_Ag- | THP, VPA, BUP, TZD, ROP, LTG |
| 23 | 50 | M | 18 | FW | Paroxysmal leg weakness | - | PTSD | CZP |
| 24 | 38 | F | 18 | FMD, FW* | Abnormal gait, right arm and leg weakness | PD_Ag- | - | BUP, DXAM/APM |
| 25 | 38 | F | 10 | FMD [†] | Paroxysmal abnormal truncal and head movements | Undiff Somatoform | - | - |
| 26 | 22 | F | 12 | PNES [†] | Motor seizures | PTSD, GAD, Ag | MDD, Social Phobia, OCD | VEN, FLX, DZP |
| 27 | 42 | F | 18 | FMD | Right arm and head tremor, dystonia (bilateral feet) | Somatoform Pain Disorder | MDD | SERT, DZP, GBP |
| 28 | 38 | F | 16 | FMD, PNES* | Paroxysmal dystonia (bilateral feet), four limb tremors, motor seizures | PD_Ag-, Social Phobia, OCD, PTSD, Somatization Disorder | MDD | CLP, LRZ, MIR, SERT, PGN, LTG |
| 29 | 49 | M | 13 | FW* [†] | Right wrist weakness | PD_Ag-, GAD | MDD | FLX, GBP, DXAM/APM |
| 30 | 50 | F | 14 | FMD, PNES | Head and bilateral arm tremor, motor seizures | MDD, PD_Ag+, PTSD, GAD, Eating Disorder NOS | Alcohol, Cocaine, Opiate and Stimulants Abuse | FLX, ZLP, LRZ |

| | | | | | | | | |
|----|----|---|----|-----------------------|--|---|---------------------------------|---------------------|
| 31 | 31 | M | 14 | PNES | Episodes of behavioral arrest with altered consciousness | PD_Ag-, PTSD | MDD, Alcohol Abuse | DLX, NRT, PZN |
| 32 | 39 | F | 16 | FW* [†] | Four limb weakness | PTSD, GAD, Somatization Disorder | MDD | CLN |
| 33 | 35 | F | 12 | PNES | Motor seizures | - | MDD | TPM, TZD |
| 34 | 54 | F | 18 | FMD | Head tremor | Dysthymia, GAD | Anorexia, PD_Ag+, Social Phobia | FLX, CLP, TZD, LXAM |
| 35 | 51 | M | 14 | FMD, FW [†] | Paroxysmal full body tremor, right arm and bilateral leg weakness | MDD, PTSD, Somatoform Pain Disorder | MDD, Social Phobia | CLP, DLX, GBP, BZP |
| 36 | 70 | M | 19 | FMD | Abnormal gait | - | Anxiety NOS | - |
| 37 | 58 | F | 18 | PNES | Motor seizures with fear and crying | Ag | MDD, Alcohol Abuse | TZD, GBP, CTP, TPM |
| 38 | 53 | F | 16 | FMD | Abnormal gait | MDD, Dysthymia, GAD, Somatoform Pain Disorder | Eating Disorder NOS | GBP, TZD, DLX |
| 39 | 49 | F | 15 | PNES | Motor seizures | PD_Ag- | MDD, PTSD | LRZ, LTG |
| 40 | 28 | F | 16 | PNES, FW [†] | Motor seizures with altered consciousness, left arm and leg weakness | Dysthymia, PDwoAG, Undiff Somatoform | PTSD, Depression NOS | GBP |
| 41 | 29 | F | 18 | PNES [†] | Motor seizures w/ altered consciousness | GAD, Specific Phobia | Eating Disorder NOS | TPM |

| | | | | | | | | |
|----|----|---|----|--------------------|--|--|------------------------------|-----------------------|
| 42 | 37 | F | 18 | PNES, FMD | Motor seizures w/ LOC, left arm tremor | PD_Ag-, Undiff Somatoform | PTSD, Social Phobia | - |
| 43 | 34 | F | 14 | FMD, FW* | Limb weakness, abnormal gait | BPAD II, Anxiety NOS | - | TZD, LRZ |
| 44 | 25 | F | 19 | FMD | Dystonia (bilateral feet) | Somatoform Pain Disorder, Anxiety NOS, BPAD II | OCD | DZP, GBP, VTX |
| 45 | 55 | F | 12 | PNES | Motor seizures w/ LOC | MDD, Dysthymia, PTSD | Social Phobia | BUP, LRZ, QTP, ZLP |
| 46 | 52 | F | 12 | FMD | Paroxysmal head and bilateral arm tremors | Depression NOS | - | SERT |
| 47 | 57 | M | 12 | FMD* | Right arm tremor | OCD, Anxiety NOS | MDD | BUP, DLX, ZLP |
| 48 | 20 | F | 14 | FMD, FW | Abnormal gait, right arm tremor and weakness | PD_Ag- | MDD | - |
| 49 | 30 | F | 18 | FMD, PNES | Motor seizures, abnormal gait | Anxiety NOS, Undiff Somatoform | - | - |
| 50 | 51 | M | 19 | PNES ⁺⁺ | Sensory symptoms and loss of time | - | MDD, Anxiety NOS | DNZ |
| 51 | 19 | M | 14 | FMD | Jerking head and limb movements | Anxiety NOS | - | - |
| 52 | 48 | F | 12 | FMD | Abnormal gait | Anxiety NOS ^{**} | Depression NOS ^{**} | AMD, SERT, LRZ |

| | | | | | | | | |
|----|----|---|----|---------------|---|--|----------------------------------|--------------------|
| 53 | 31 | M | 14 | PNES | Minor motor seizures | PD_Ag-, Undiff Somatoform | PTSD, ETOH Abuse | PGN, PZN, DLX |
| 54 | 20 | F | 12 | PNES | Episodes of behavioral arrest with altered consciousness | Somatization Disorder, Anxiety NOS | Depression NOS | BSP, ECP, LRZ, TZD |
| 55 | 64 | F | 15 | FW* | Generalized weakness | PTSD, OCD, Somatization Disorder, Depression NOS | MDE, PD_Ag+, Eating Disorder NOS | DLX, MIR |
| 56 | 29 | F | 18 | FMD, FW, PNES | Minor motor seizures with altered consciousness, abnormal gait, left arm and bilateral leg weakness | Somatization Disorder, Anxiety NOS | - | GBP, LRZ |

†Indicates that these patients had splitting of the midline functional numbness. ††Indicates that patient also had a functional cognitive disorder. *Indicates that these patients also had functional voice symptoms. **Psychiatric diagnoses made by chart review, as structured psychiatric interview not performed. M, male; F, female; NES, Nonepileptic Seizures; FW, Functional Weakness; FMD, Functional Movement Disorders; NOS, not otherwise specified; MDD, Major Depressive Disorder; PD_Ag+, Panic Disorder with Agoraphobia; PD_Ag-, Panic Disorder without Agoraphobia; GAD, Generalized Anxiety Disorder; Undiff Somatoform, Undifferentiated Somatoform Disorder; PTSD, Post-Traumatic Stress Disorder; OCD, Obsessive Compulsive Disorder; Ag, Agoraphobia without Panic Disorder; LOC, Loss of Consciousness; AMD,

Amantadine; APM, Amphetamine; APZ, Aripiprazole; ATP, Amitriptyline; BSP, Buspirone; BUP, Bupropion; BZP, Benzotropine; CLN, Clonidine; CLP, Clonazepam; CTP, Citalopram; CZP, Carbamazepine; DLX, Duloxetine; DNZ, donepezil; DOX, Doxepin; DVX, Desvenlafaxine; DXAM, Dextroamphetamine; DZP, Diazepam; ECP, Escitalopram; FLX, Fluoxetine; GBP, Gabapentin; LRZ, Lorazepam; LSD, Lysergic Acid Diethylamide; LTG, Lamotrigine; LXAM, Lisdexamfetamine; MIR, Mirtazapine; NRT, Nortriptyline; PGN, Pregabalin; PZN, Prazosin; QTP, Quetiapine; ROP, Ropinirole; SERT, Sertraline; THP, Trihexyphenidyl; TPM, Topiramate; TZD, Trazodone; VEN, Venlafaxine; VPA, Valproic Acid; VTX, Vortioxetine; ZLP, Zolpidem.

Supplemental Table 2. Demographic characteristics and within group univariate associations with self-reported attachment styles (RSQ) in motor functional neurological disorders.

| | | RSQ Secure | RSQ Fearful | RSQ Preoccupied | RSQ Dismissing |
|--|-----------|-------------------------|-------------------------|-------------------------|-------------------------|
| | Mean±SD | Test Statistic; p-value | Test Statistic; p-value | Test Statistic; p-value | Test Statistic; p-value |
| Age | 40.2±13.0 | 0.15; 0.28 | -0.02; 0.91 | -0.08; 0.55 | 0.10; 0.48 |
| Female (41) v. Male (15) | - | 0.31; 0.76 | 1.01; 0.32 | -0.27; 0.79 | -1.17; 0.25 |
| White (51) v. Non-white (5) | - | 0.74; 0.47 | 1.21; 0.23 | -0.48; 0.63 | -2.13; 0.04 |
| College (21) v. No College (35) | - | 1.46; 0.15 | 1.27; 0.21 | 1.45; 0.15 | 1.28; 0.21 |
| Illness Duration in years | 3.3±3.7 | -0.02; 0.91 | -0.04; 0.78 | 0.17; 0.21 | -0.12; 0.39 |
| Married (23) v Not Married (33) | - | 1.35; 0.18 | -2.09; 0.04 | -1.23; 0.22 | -2.41; 0.02 |
| Disabled (25) v Not Disabled (31) | - | 0.99; 0.33 | -0.62; 0.54 | -1.17; 0.25 | 0.02; 0.99 |
| Full Time Employed or Student (21) v. Other (35) | - | 1.12; 0.27 | 0.67; 0.51 | 1.03; 0.31 | 0.16; 0.88 |
| PNES (24) v. Other Motor FND | - | 1.09; 0.28 | 0.04; 0.97 | 1.51; 0.14 | -0.48; 0.63 |
| FMD (29) v. Other Motor FND | - | 2.27; 0.03 | -0.65; 0.52 | -0.81; 0.42 | 0.05; 0.96 |
| FW (21) v. Other Motor FND | - | 1.03; 0.31 | 1.52; 0.13 | 0.48; 0.64 | 1.08; 0.28 |
| SOMS:CD | 8.4±7.2 | 0.18; 0.19 | 0.15; 0.27 | 0.03; 0.83 | 0.06; 0.65 |
| PHQ-15 | 13.2±5.0 | -0.05; 0.74 | 0.35; 0.009 | 0.15; 0.27 | -0.03; 0.82 |
| SDQ-20 | 32.2±10.1 | 0.15; 0.28 | 0.43; 0.001 | 0.08; 0.58 | 0.06; 0.68 |
| SF-36 Mental | 47.9±22.9 | 0.30; 0.03 | -0.48; <0.001 | -0.18; 0.18 | -0.04; 0.75 |
| SF-36 Physical | 46.4±23.1 | -0.14; 0.31 | -0.23; 0.09 | 0.02; 0.90 | -0.05; 0.71 |
| DES | 19.2±14.4 | -0.06; 0.64 | 0.57; <0.001 | 0.38; 0.004 | 0.11; 0.40 |
| PTSD-CL5* | 26.7±20.3 | -0.30; 0.03 | 0.49; <0.001 | 0.07; 0.63 | 0.12; 0.39 |
| LEC 'Happened to Me' | 4.2±2.8 | -0.11; 0.41 | 0.37; 0.005 | 0.05; 0.70 | 0.25; 0.06 |
| CTQ-Abuse | 28.5±13.6 | -0.23; 0.09 | 0.55; <0.001 | 0.23; 0.09 | 0.02; 0.87 |
| CTQ-Neglect | 20.3±8.8 | -0.09; 0.52 | 0.34; 0.01 | 0.14; 0.30 | -0.13; 0.33 |
| BDI-II | 18.7±12.5 | -0.20; 0.14 | 0.61; <0.001 | 0.17; 0.22 | 0.16; 0.24 |
| STAI-T | 85.8±23.8 | -0.42; 0.001 | 0.56; <0.001 | 0.28; 0.04 | 0.17; 0.23 |
| CD-RISC | 64.0±16.5 | 0.47; <0.001 | -0.30; 0.03 | -0.22; 0.11 | 0.07; 0.59 |
| TAS | 56.1±13.2 | -0.47; <0.001 | 0.38; 0.004 | 0.20; 0.14 | 0.11; 0.44 |
| NEO-Neuroticism | 25.4±9.0 | -0.44; 0.001 | 0.48; <0.001 | 0.34; 0.01 | 0.14; 0.30 |

| | | | | | |
|-----------------------|-----------|------------------------|------------------------|--------------------|-------------|
| NEO-Extraversion | 24.1±7.4 | 0.47; <0.001 | -0.40; 0.003 | -0.16; 0.23 | -0.13; 0.36 |
| NEO-Openness | 26.1±5.7 | 0.16; 0.25 | 0.04; 0.76 | -0.12; 0.36 | 0.25; 0.06 |
| NEO-Agreeableness | 35.5±5.0 | -0.10; 0.47 | 0.21; 0.12 | -0.09; 0.49 | 0.13; 0.34 |
| NEO-Conscientiousness | 31.8±7.3 | 0.10; 0.45 | -0.08; 0.55 | -0.32; 0.02 | 0.15; 0.28 |
| BIS | 64.2±10.8 | -0.01; 0.92 | 0.15; 0.27 | 0.28; 0.04 | -0.11; 0.43 |
| PCS* | 16.5±12.2 | -0.23; 0.09 | 0.51; <0.001 | 0.32; 0.02 | 0.05; 0.74 |

Test statistic refers to independent sample t-test for dichotomous group comparisons or Pearson correlation coefficient for continuous variables. *Indicates that one subject had missing data for the PTSD-CL5 and another subject had missing data for the PCS. RSQ, Relationship Scales Questionnaire; FND, Functional Neurological Disorder; PNES, psychogenic non-epileptic seizures; FMD, functional movement disorder; FW, functional weakness; SD, standard deviation; SOMS: CD, Conversion Disorder Subscale of the Screening for Somatoform Symptoms-7 scale; PHQ-15, Patient Health Questionnaire-15; SDQ-20, Somatoform Dissociation Questionnaire-20; SF-36; Short Form Health Survey-36; DES, Dissociative Experience Scale; PTSD-CL5, Post-Traumatic Stress Disorder Checklist for DSM-5; LEC, Life Events Checklist; CTQ, Childhood Trauma Questionnaire; BDI-II, Beck Depression Inventory-II; STAI-T, Spielberger State Trait Anxiety Inventory Total Score; CD-RISC, Connor-Davidson Resilience Scale; TAS, Toronto Alexithymia Score; NEO, NEO Five-Factor Inventory-3; BI, Barrett Impulsivity Scale; PCS, Pain Catastrophizing Scale.