

Association of Glutathione s-transferase M1 and T1 gene polymorphisms with the susceptibility to acquired sensorineural hearing loss: a systematic review and meta-analysis

Shimin Zong¹⁺, Xue Zeng²⁺, Yexiao Guan¹⁺, Tianyi Liu¹, Pan Luo¹, Fangmin Wan¹, Yanji Qu¹, Pei Chen³, Hongjun Xiao^{1*}

¹ Department of Otorhinolaryngology, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan 430022, China

²Department of Obstetrics and Gynecology, Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan 430032, China

³Department of Otorhinolaryngology, Wuhan Integrated TCM and Western Medicine Hospital (Wuhan No. 1 hospital), Tongji Medical College, Huazhong University of Science and Technology, Wuhan 430022, China

*Corresponding. xhjent_whxh@hust.edu.cn (Hongjun Xiao)

⁺The authors contribute equally to this work.

	Gene name	Before meta-regression	After meta-regression	Variation
Tau - squared	GST M1	0.0901	0.06719	25.4%
	GST T1	0.1757	0.2011	-

Supplementary Table 1. Results of meta-regression considering sample size, publication year and NOS scores of the included studies. GST: glutathione s-transferase.