

Supporting information (S)

Biological activities of *Ficus carica* latex for potential therapeutics in Human Papillomavirus (HPV) related cervical cancers

Arshia Ghanbari ¹, Adam Le Gresley ², Declan Naughton ², Nikolai Kuhnert ³, Diana Sirbu ³, G. Hossein Ashrafi ^{1*}


¹ Kingston University London, Cancer theme, School of Life Science, Pharmacy and Chemistry, SEC Faculty, Kingston upon Thames, KT1 2EE, London, UK

² Kingston University London, School of Life Science, Pharmacy and Chemistry, SEC Faculty, Kingston upon Thames, KT1 2EE, London, UK

³ Department of Life Sciences & Chemistry Faculty of Health, Jacobs University Bremen, Campusring 8, 28759 Bremen, Germany


* Corresponding Author: H.ashrafi@kingston.ac.uk

Supporting information 1 (S1)




S1 ^1H NMR Spectra in $\text{D}_2\text{O}/\text{H}_2\text{O}$ 90:10% of Fig latex extracts. Top – Fig Latex Gum, Middle, Fig Latex Powder, Bottom, Fig Latex Supernatant.

Supporting information 2 (S2)



S2 ¹H NMR Aromatic region of petroleum ether extract of fig latex supernatant showing at least two aromatic components, suggested asferrulic, caffeic or chlorogenic acids.

Supporting information 3 (S3)




S3. Full unprocessed original scanned gels for western blots shown in figures 8.

M= Molecular weight marker

- = untreated

+ = Treated with Fig latex

Supporting information 4 (S4)



S4. Expression of p53 in HaCaT cells

Full unprocessed western blot.

M=Molecular weight marker