

Supplementary Table 1. Demographic and clinical characteristics of the patients at baseline.

	Exenatide (n=23)	Placebo (n=22)	p-value
Age (year) – mean ± SD [range]	37.1 ± 10.6 [19-65]	34.5 ± 10.1 [19-56]	0.40
Sex (male / female) – no. (%)	11 (48) / 12 (52)	10 (45) / 12 (55)	0.87
Ethnicity (caucasian / mongolian) – no. (%)	21 (91) / 2 (9)	19 (86) / 3 (14)	0.60
Illness duration (weeks) – mean ± SD [range]	689 ± 463 [38-1404]#	578 ± 426 [26-1404]	0.41
Education (years) – mean ± SD [range]	12.6 ± 2.9 [8-18]	12.1 ± 2.6 [7-20]	0.51
Current smoker – no. (%)	7 (30)	1 (5)	0.02
Body weight (kg) – mean ± SD [range]	117.1 ± 16.7 [84.4-150.7]	110.6 ± 17.7 [88.2-149.7]	0.21
BMI (kg/m²) – mean ± SD [range]	39.2 ± 3.8 [31.0-48.0]	38.4 ± 6.1 [30.1-55.1]	0.59
Diagnosis			
Schizophrenia, F20.X – no. (%)	21 (91)	20 (91)	-
Schizoaffective disorders, F25.X – no. (%)	2 (9)	2 (9)	
Antipsychotic medication			
Typical monotherapy – no. (%)	1 (4)	1 (5)	0.97
Atypical monotherapy – no. (%)	15 (65)	11 (50)	0.30
Polypharmacy – no. (%)	7 (30)	10 (45)	0.30

One observation missing.