

Supplementary figure 1 for Kovács et al. “Cadaverine, a metabolite of the microbiome, reduces breast cancer aggressiveness through trace amino acid receptors”

Tünde Kovács¹, Edit Mikó^{1,5}, András Vida^{1,5}, Éva Sebő⁶, Judit Toth³, Tamás Csonka⁴, Anita Boratkó¹, Gyula Ujlaki¹, Gréta Lente¹, Patrik Kovács¹, Dezső Tóth³, Péter Árkosy³, Borbála Kiss², Gábor Méhes⁴, James J. Goedert⁷, Péter Bai^{1,5,8,*}

Departments of ¹Medical Chemistry, ²Dermatology, ³Oncology and ⁴Pathology Faculty of Medicine, University of Debrecen, 4032, Hungary;

⁵MTA-DE Lendület Laboratory of Cellular Metabolism, Debrecen, 4032, Hungary;

⁶Kenézy Breast Center, Kenézy Gyula County Hospital, Debrecen, 4032, Hungary;

⁷National Cancer Institute, National Institutes of Health, Bethesda, 20982 MD, USA



⁸Research Center for Molecular Medicine, Faculty of Medicine, University of Debrecen, 4032, Hungary;

Running title: Cadaverine production modulates breast cancer

*Whom correspondence should be sent to:



Peter Bai, PhD, DSc University of Debrecen, Department of Medical Chemistry, 4032 Debrecen, Egyetem tér 1., Hungary, Tel. +36 52 412 345; Fax. +36 52 412 566, e-mail: baip@med.unideb.hu

A




mesenchymal
epithelial


D



B



C



E

