

Supplemental Table 2. Multivariable logistic regression of nailfold capillary blood speed, diameter, and flow in relation to primary open-angle glaucoma (POAG) patients not using topical beta-blocker medication (n=26) or using topical beta-blocker medication (n=36) vs. controls (n=57)

| Exposure (covariates indented) | POAG not using topical beta-blockers vs. Controls | | POAG using topical beta-blocker in at least one eye vs. Controls | |
|---|---|---------------|--|-------------------|
| | Odds ratio (95% confidence interval) | P value | Odds ratio (95% confidence interval) | P value |
| Mean speed in microns per second | 0.994 (0.990, 0.997) | 0.0013 | 0.993 (0.990, 0.996) | <0.0001 |
| Body mass index in kg/m ² | 0.99 (0.83, 1.17) | 0.95 | 0.97 (0.84, 1.11) | 0.68 |
| Mean arterial pressure in mmHg | 0.99 (0.93, 1.06) | 0.75 | 0.98 (0.93, 1.03) | 0.45 |
| Pulse in beats per minute | 1.10 (1.02, 1.19) | 0.015 | 1.03 (0.98, 1.08) | 0.33 |
| Female sex | 0.59 (0.091, 3.49) | 0.56 | 0.59 (0.17, 1.92) | 0.39 |
| Age in years | 1.00 (0.93, 1.08) | 0.98 | 1.00 (0.94, 1.08) | 0.97 |
| Caucasian race | 2.91 (0.32, 30.7) | 0.34 | 3.60 (0.69, 22.8) | 0.14 |
| Connective tissue disease | 0.49 (0.069, 2.77) | 0.44 | 0.81 (0.20, 3.20) | 0.76 |
| Diabetes mellitus | 0.074 (0.0017, 1.55) | 0.12 | 0.17 (0.013, 1.31) | 0.12 |
| Use of blood thinning medication | 0.72 (0.11, 4.95) | 0.73 | 0.68 (0.18, 2.44) | 0.55 |
| Use of blood pressure medication | 0.20 (0.020, 1.45) | 0.13 | 1.22 (0.30, 5.33) | 0.78 |
| Ever smoker (vs. never) | 2.03 (0.51, 8.66) | 0.32 | 1.08 (0.30, 3.87) | 0.90 |
| Mean diameter in microns | 0.67 (0.42, 1.01) | 0.065 | 0.74 (0.51, 1.06) | 0.11 |
| Body mass index in kg/m ² | 0.90 (0.77, 1.03) | 0.15 | 0.91 (0.81, 1.02) | 0.12 |
| Mean arterial pressure in mmHg | 1.00 (0.95, 1.06) | 0.95 | 1.01 (0.97, 1.05) | 0.67 |
| Pulse in beats per minute | 1.08 (1.02, 1.15) | 0.018 | 1.01 (0.97, 1.06) | 0.62 |
| Female sex | 0.61 (0.12, 3.08) | 0.54 | 0.73 (0.26, 2.04) | 0.55 |
| Age in years | 0.98 (0.92, 1.05) | 0.58 | 0.98 (0.93, 1.05) | 0.59 |
| Caucasian race | 4.20 (0.52, 37.8) | 0.17 | 3.59 (0.80, 18.9) | 0.11 |
| Connective tissue disease | 0.31 (0.047, 1.59) | 0.18 | 1.11 (0.33, 3.66) | 0.86 |
| Diabetes mellitus | 0.13 (0.0033, 2.01) | 0.19 | 0.24 (0.025, 1.44) | 0.15 |
| Use of blood thinning medication | 0.63 (0.11, 3.52) | 0.60 | 0.78 (0.24, 2.46) | 0.68 |
| Use of blood pressure medication | 0.44 (0.064, 2.69) | 0.38 | 1.55 (0.45, 5.50) | 0.49 |
| Ever smoker (vs. never) | 3.14 (0.90, 12.4) | 0.084 | 1.39 (0.45, 4.30) | 0.56 |
| Mean flow in picoliters per second | 0.94 (0.90, 0.97) | 0.0017 | 0.93 (0.90, 0.96) | <0.0001 |
| Body mass index in kg/m ² | 0.96 (0.81, 1.13) | 0.67 | 0.96 (0.83, 1.09) | 0.53 |
| Mean arterial pressure in mmHg | 0.99 (0.93, 1.06) | 0.82 | 0.98 (0.93, 1.03) | 0.41 |
| Pulse in beats per minute | 1.10 (1.03, 1.20) | 0.0095 | 1.03 (0.98, 1.09) | 0.23 |
| Female sex | 0.31 (0.042, 1.87) | 0.21 | 0.54 (0.15, 1.80) | 0.33 |
| Age in years | 0.98 (0.91, 1.06) | 0.62 | 0.99 (0.93, 1.06) | 0.73 |
| Caucasian race | 3.54 (0.37, 40.0) | 0.28 | 4.58 (0.84, 31.6) | 0.093 |
| Connective tissue disease | 0.41 (0.055, 2.39) | 0.35 | 0.87 (0.21, 3.50) | 0.84 |
| Diabetes mellitus | 0.047 (0.00091, 1.13) | 0.084 | 0.11 (0.0069, 1.02) | 0.078 |
| Use of blood thinning medication | 0.43 (0.061, 2.86) | 0.38 | 0.65 (0.17, 2.41) | 0.52 |
| Use of blood pressure medication | 0.35 (0.039, 2.63) | 0.32 | 1.67 (0.40, 7.55) | 0.48 |
| Ever smoker (vs. never) | 2.82 (0.70, 12.9) | 0.16 | 1.50 (0.40, 5.72) | 0.55 |