

Supplemental Results:

Table S1. Additional BP Outcomes in the Intervention and Control Groups[†]

	Intervention		Control		ANOVA <i>P</i> -Values		
	Pre-Study	During Study	Pre-Study	During Study	Group Effect	Time Effect	Inter-action Effect
Average Highest SBP (mm Hg)	155 ± 21	153 ± 15	174 ± 17	172 ± 13	<0.0001*	0.41	0.94
Average Lowest SBP (mm Hg)	111 ± 15	108 ± 11	118 ± 14	122 ± 16	0.0006*	0.82	0.26
Max. Value of the Highest SBP (mm Hg)	188 ± 27	192 ± 25	207 ± 21	206 ± 25	0.01*	0.77	0.64
Min. Value of the Lowest SBP (mm Hg)	86 ± 19	83 ± 16	92 ± 15	90 ± 16	0.07	0.52	0.93
Average Highest MAP (mm Hg)	102 ± 12	101 ± 7	113 ± 13	112 ± 11	<0.0001*	0.52	0.90
Average Lowest MAP (mm Hg)	77 ± 10	75 ± 8	81 ± 9	84 ± 11	0.0003*	0.92	0.18
Max. Value of the Highest MAP (mm Hg)	130 ± 17	132 ± 18	135 ± 22	138 ± 20	0.12	0.50	0.86
Min. Value of the Lowest MAP (mm Hg)	60 ± 12	60 ± 10	62 ± 9	63 ± 9	0.35	0.85	0.73

[†]Data are presented as mean ± SD; SBP, systolic blood pressure; MAP, mean arterial pressure. “Average”, “highest”, and “lowest” values refer to over the 25 during study dialysis sessions for each patient.

**p* < 0.05