

Impact of gold nanoparticles shape on their cytotoxicity against human osteoblast and osteosarcoma in *in vitro* model. Evaluation of the safety of use and anti-cancer potential.

Karol P. Steckiewicz¹, Ewelina Barcińska¹, Anna Malankowska², Agata Zauszkiewicz – Pawlak⁴, Grzegorz Nowaczyk³, Adriana Zaleska-Medynska², Iwona Inkielewicz-Stępnik¹

¹Department of Medical Chemistry, Medical University of Gdańsk, Dębinki 1, 80-211 Gdansk, Poland

²Department of Environmental Technology, Faculty of Chemistry, University of Gdansk, Wita Stwosza 63, 80-308 Gdansk, Poland

³NanoBioMedical Center, Adam Mickiewicz University, 61-614 Poznan, Poland

⁴Department of Histology, Medical University of Gdańsk, Dębinki 1, 80-211 Gdansk, Poland

Corresponding author: Iwona Inkielewicz-Stępnik, iinkiel@gumed.edu.pl, tel. +48 58 349 1450

Electronic Supplementary Material




Fig. Log(concentration) vs. cell viability plots. Non-linear regression analysis: log(inhibitor) vs. normalized response has been performed to calculate log IC₅₀ values. Calculation were made for three cell lines (hFOB1.19, MG-63 and 143B), using (a) MTT and (b) NR assay. The obtained data are reported as the mean ± SD for triplicate determination of 3 separate experiments