

Supporting information

Selective Capture and Purification of MicroRNAs and Intracellular Proteins through Antisense-vectorized Magnetic Nanobeads

Isabel Gessner¹, Xiaojie Yu^{2,3}, Christian Jüngst⁴, Annika Klimpel⁵, Lingyu Wang^{2,3}, Thomas Fischer¹, Ines Neundorff⁵, Astrid Schauss⁴, Margarete Odenthal^{2,3,6}, Sanjay Mathur^{1,*}

¹ Institute of Inorganic Chemistry, University of Cologne, Greinstr. 6, 50939 Cologne, Germany

² Institute for Pathology, University Hospital of Cologne, Kerpener Str. 62, 50924 Cologne, Germany

³ Center for Molecular Medicine (CMMC), University of Cologne, Robert-Koch-Straße 21, 50931 Cologne, Germany

⁴ Cluster of Excellence - Cellular Stress Responses in Aging-Associated Diseases (CECAD), Imaging Facility, University of Cologne, Joseph-Stelzmann-Str. 26, 50931 Cologne, Germany

⁵ Institute of Biochemistry, University of Cologne, Zulpicher Str. 47, 50674 Cologne, Germany

⁶ Center of Integrative Oncology, University Clinic of Cologne and Bonn, Germany

*corresponding author: sanjay.mathur@uni-koeln.de




Figure S 1: Orthogonal view of confocal microscopic stackings performed for scramble functionalized magnetic beads in Huh7 cells.




Figure S 2: Confocal microscopic images of Huh7 cells incubated with scramble functionalized magnetic beads for incubation times between 0-24 hours. The cytoskeleton was stained with phalloidin green, while the nucleus was stained with DAPI. The scale bar in all images refers to 25 μ m.