

Supplementary Table S-1. Showing number of subjects with missing values for each clinical and serological measurement for the controls, SLE patients and the two suggested subgroups of aPL+ SLE and SSA/SSB+ SLE.

Missing Values	Controls	SLE	aPL+ SLE	SSA/SSB+ SLE
	n=316	n=378	n=66	n=63
Sampling Age	0	0	0	0
Gender	0	0	0	0
Nephritis	3	0	0	0
SLEDAI-2K	316	0	0	0
SLAM	316	0	0	0
SLICC.total	316	0	0	0
Disease Duration	316	0	0	0
Acetylsalicylic Acid	313	376	66	63
Azathioprin	316	327	55	58
Calcium and D supplementation	313	343	60	58
Cyclophosphamide	316	377	66	63
D vitamin	2	9	0	2
Esomeprazol	314	368	64	62
Folic Acid	315	354	60	61
Furosemide	315	352	62	62
Hyrdoxychloroquine	316	311	57	51
Ibuprofen	312	365	62	60
Karbamazepin	316	374	66	62
Levothyroxine	296	335	60	55
Methotrexate	316	367	64	63
Mycophenolate	316	351	63	60
Paracetamol	304	319	59	54
Rituximab	314	69	11	8
SNRI	0	0	0	0
SSRI	0	0	0	0
Warfarin	315	339	45	58
A1-antitrypsin	4	7	3	0
Apo A	13	76	13	9
Apo B	13	76	13	9
C1q	316	22	2	2
C2	316	43	9	5
C3	0	9	2	1
C3dg	316	121	24	17
C4	0	9	2	1
TCC	0	0	0	0
Creatinine	0	0	0	0
SLEDAI-2K Proteinurea =4	316	0	0	0
ESR	8	15	3	3
Fibrinogen	8	11	3	0
Homocystein	13	76	13	9
Hs CRP	11	2	0	0
IgA	4	0	0	0
IgG	4	0	0	0
IgM	4	0	0	0
Rf-IgM	95	29	8	5

Rf-IgA	97	53	7	19
Rf-IgG	112	53	8	10
Leucocytes	0	1	1	0
Lymphocytes	0	3	2	0
Neutrophils	0	3	2	0
Platelets	0	1	1	0
TG	0	1	0	0
VCAM	6	76	12	10
