

# **Optical manipulation of sphingolipid biosynthesis using photoswitchable ceramides**

Matthijs Kol<sup>\$1</sup>, Ben Williams<sup>\$2</sup>, Henry Toombs-Ruane<sup>2</sup>, Henri G. Franquelim<sup>3</sup>, Sergei Korneev<sup>1</sup>, Christian Schroer<sup>1</sup>, Petra Schwille<sup>3</sup>, Dirk Trauner<sup>\*2,4</sup>, Joost C.M. Holthuis<sup>\*1,5</sup>, James Allen Frank<sup>\*2,6</sup>

<sup>1</sup>Molecular Cell Biology Division, Department of Biology/Chemistry, University of Osnabrück, 49076 Osnabrück, Germany. <sup>2</sup>Department of Chemistry and Center for Integrated Protein Science, Ludwig Maximilians University Munich, Butenandtstraße 5-13, 81377 Munich, Germany. <sup>3</sup>Department of Cellular and Molecular Biophysics, Max Planck Institute of Biochemistry, Am Klopferspitz 18, 82152 Martinsried, Germany. <sup>4</sup>Department of Chemistry, New York University, 100 Washington Square East, New York, NY 10003-6699, USA. <sup>5</sup>Membrane Biochemistry & Biophysics, Bijvoet Center and Institute of Biomembranes, Utrecht University, 3584 CH Utrecht, the Netherlands. <sup>6</sup>*present address:* The Vollum Institute, Oregon Health & Science University, 3181 S.W. Sam Jackson Park Rd., Portland, Oregon 97239, USA

<sup>\$</sup>these authors contributed equally to this work

\*co-corresponding authors


James Allen Frank – frankja@ohsu.edu

Joost C.M. Holthuis - joost.holthuis@biologie.uni-osnabrueck.de


Dirk Trauner – dirktrauner@nyu.edu

## **Supplementary File 1 - NMR Spectra**


<sup>1</sup>H NMR of 1-(but-3-yn-1-yl)-4-nitrobenzene (2)




<sup>13</sup>C NMR of 1-(but-3-yn-1-yl)-4-nitrobenzene (2)




<sup>1</sup>H NMR of 4-(but-3-yn-1-yl)aniline (3)




<sup>13</sup>C NMR of 4-(but-3-yn-1-yl)aniline (3)




<sup>1</sup>H NMR of 4-(4-((4-(but-3-yn-1-yl)phenyl)diazenyl)phenyl)butanoic acid (cFAAzo-4)




<sup>13</sup>C NMR of 4-(4-((4-(but-3-yn-1-yl)phenyl)diazenyl)phenyl)butanoic acid (cFAAzo-4)




<sup>1</sup>H NMR of 4-(4-(but-3-yn-1-yl)phenyl)diazenylphenyl)-N-((2S,3R,E)-1,3-dihydroxyoctadec-4-en-2-yl)butanamide (caCer-1)




<sup>13</sup>C NMR of 4-(4-(but-3-yn-1-yl)phenyl)diazenylphenyl)-N-((2S,3R,E)-1,3-dihydroxyoctadec-4-en-2-yl)butanamide (caCer-1)




<sup>1</sup>H NMR of 4-((4-(but-3-yn-1-yl)phenyl)diaz恒)benzoic acid (cFAAzo-1)




<sup>13</sup>C NMR of 4-((4-(but-3-yn-1-yl)phenyl)diaz恒)benzoic acid (cFAAzo-1)




<sup>1</sup>H NMR of 4-(4-(but-3-yn-1-yl)phenyl)diazenyl-N-((2*S*,3*R*,*E*)-1,3-dihydroxyoctadec-4-en-2-yl)benzamide (caCer-2)




<sup>13</sup>C NMR of 4-(4-(but-3-yn-1-yl)phenyl)diazenyl-N-((2*S*,3*R*,*E*)-1,3-dihydroxyoctadec-4-en-2-yl)benzamide (caCer-2)




<sup>1</sup>H NMR of pentadec-7-yn-1-ol (S1)




<sup>13</sup>C NMR of pentadec-7-yn-1-ol (S1)




<sup>1</sup>H NMR of pentadec-14-yn-1-ol (**S2**)




<sup>1</sup>H NMR of pentadec-14-ynoic acid (**4**)




<sup>1</sup>H NMR of 4-((4-propylphenyl)diazenyl)benzyl alcohol (5)




<sup>13</sup>C NMR of 4-((4-propylphenyl)diazenyl)benzyl alcohol (5)




<sup>1</sup>H NMR of 4-((4-propylphenyl)diazenyl)benzaldehyde (6)




<sup>13</sup>C NMR of 4-((4-propylphenyl)diazenyl)benzaldehyde (6)




<sup>1</sup>H NMR of 1-(4-propylphenyl)-2-(4-vinylphenyl)diazene (7)




<sup>13</sup>C NMR of 1-(4-propylphenyl)-2-(4-vinylphenyl)diazene (7)




<sup>1</sup>H NMR of 4-iodo-4'-methylazobenzene (S3)




<sup>13</sup>C NMR of 4-iodo-4'-methylazobenzene (S3)




<sup>1</sup>H NMR of 3-(4-(*p*-tolylidazeny)phenyl)propanal (**S4**)




<sup>13</sup>C NMR of 3-(4-(*p*-tolylidazeny)phenyl)propanal (**S4**)




<sup>1</sup>H NMR of **1-(4-(but-3-en-1-yl)phenyl)-2-(*p*-tolyl)diazene (S5)**




<sup>13</sup>C NMR of **1-(4-(but-3-en-1-yl)phenyl)-2-(*p*-tolyl)diazene (S5)**




<sup>1</sup>H NMR of *N*-Boc-*(R)*-1-((*S*)-2,2-dimethyloxazolidin-4-yl)prop-2-en-1-ol (8)




<sup>13</sup>C NMR of *N*-Boc-*(R)*-1-((*S*)-2,2-dimethyloxazolidin-4-yl)prop-2-en-1-ol (8)




<sup>1</sup>H NMR of **N-Boc-(R,E)-1-((S)-2,2-dimethyloxazolidin-4-yl)-3-((4-(4-propylphenyl)diazenyl)phenyl)prop-2-en-1-ol (9)**




<sup>13</sup>C NMR of **N-Boc-(R,E)-1-((S)-2,2-dimethyloxazolidin-4-yl)-3-((4-(4-propylphenyl)diazenyl)phenyl)prop-2-en-1-ol (9)**




<sup>1</sup>H NMR of (*2S,3R,E*)-2-amino-5-(4-((4-propylphenyl)diazenyl)phenyl) pent-4-ene-1,3-diol (aSph-1)




<sup>13</sup>C NMR of (*2S,3R,E*)-2-amino-5-(4-((4-propylphenyl)diazenyl)phenyl) pent-4-ene-1,3-diol (aSph-1)




<sup>1</sup>H NMR of *N*-(*2S,3R,E*)-1,3-dihydroxy-5-(4-((4-propylphenyl) diazenyl)phenyl)pent-4-en-2-yl)pentadec-14-ynameide (caCer-3)




<sup>13</sup>C NMR of *N*-(*2S,3R,E*)-1,3-dihydroxy-5-(4-((4-propylphenyl) diazenyl)phenyl)pent-4-en-2-yl)pentadec-14-ynameide (caCer-3)




<sup>1</sup>H NMR of compound **N-Boc-(R,E)-1-((S)-2,2-dimethyloxazolidin-4-yl)-6-(4-(p-tolyldiazenyl)phenyl)hex-2-en-1-ol (S6)**




<sup>13</sup>C NMR of compound **N-Boc-(R,E)-1-((S)-2,2-dimethyloxazolidin-4-yl)-6-(4-(p-tolyldiazenyl)phenyl)hex-2-en-1-ol (S6)**




<sup>1</sup>H NMR of (*2S,3R,E*)-2-amino-7-(4-(*p*-tolylidazenyl)phenyl)hept-4-ene-1,3-diol (aSph-2)



<sup>13</sup>C NMR of (*2S,3R,E*)-2-amino-7-(4-(*p*-tolylidazenyl)phenyl)hept-4-ene-1,3-diol (aSph-2)



<sup>1</sup>H NMR of *N*-(2*S*,3*R*,*E*)-1,3-dihydroxy-7-(4-(*p*-tolylidazenyl) phenyl)hept-4-en-2-yl)pentadec-14-ynamide (caCer-4)



<sup>13</sup>C NMR of *N*-(2*S*,3*R*,*E*)-1,3-dihydroxy-7-(4-(*p*-tolylidazenyl) phenyl)hept-4-en-2-yl)pentadec-14-ynamide (caCer-4)

