

Supplementary Table 4. Urinary concentrations of BPA, BPF, and BPS according to population characteristics

Variable	No.	BPA (ng/mL)		BPF (ng/mL)		BPS (ng/mL)	
		Median (IQR)	<i>P</i> value	Median (IQR)	<i>P</i> value	Median (IQR)	<i>P</i> value
Total	745	1.23 (0.63–2.36)		0.30 (0.14–0.93)		0.28 (0.12–0.66)	
Sex			0.83		0.20		0.39
Men	371	1.25 (0.63–2.37)		0.26 (0.14–0.94)		0.27 (0.11–0.59)	
Women	374	1.22 (0.63–2.25)		0.34 (0.02–0.18)		0.29 (0.13–0.70)	
Race/ethnicity			0.17		0.26		0.19
Hispanic	253	1.10 (0.61–2.27)		0.23 (0.14–0.59)		0.38 (0.14–0.94)	
Non-Hispanic white	190	1.24 (0.59–2.14)		0.39 (0.14–1.17)		0.22 (0.11–0.56)	
Non-Hispanic black	191	1.80 (0.99–3.09)		0.36 (0.14–0.93)		0.36 (0.17–0.11)	
Other	111	1.04 (0.58–2.08)		0.14 (0.14–0.43)		0.28 (0.11–0.66)	
Family income to poverty ratio			0.33		0.36		0.37
≤1.30	350	1.29 (0.74–2.46)		0.23 (0.14–0.72)		0.30 (0.10–0.68)	
1.31–3.49	203	1.32 (0.64–2.51)		0.26 (0.14–0.97)		0.24 (0.12–0.61)	
>3.50	135	1.10 (0.59–2.02)		0.43 (0.14–1.13)		0.26 (0.13–0.60)	
Missing	57	0.82 (0.42–1.49)		0.21 (0.14–0.65)		0.41 (0.16–0.93)	
TV watching			0.16		0.45		0.45
<2 hr/day	279	1.33 (0.77–2.40)		0.34 (0.14–0.99)		0.28 (0.13–0.63)	
≥2 hr/day	466	1.12 (0.59–2.32)		0.28 (0.14–0.87)		0.27 (0.11–0.66)	
General obesity			0.59		0.53		0.39
Yes	165	1.34 (0.76–2.45)		0.32 (0.14–1.08)		0.30 (0.13–0.74)	
No	580	1.21 (0.60–2.31)		0.30 (0.14–0.90)		0.27 (0.12–0.62)	
Abdominal obesity			0.64		0.72		0.58
Yes	262	1.28 (0.63–2.55)		0.29 (0.14–1.04)		0.28 (0.11–0.70)	
No	468	1.21 (0.63–2.31)		0.30 (0.14–0.86)		0.27 (0.12–0.62)	

All the variables were adjusted using population weights for the sample in which BPA concentrations were measured except number (unweighted sample size).

BPA, bisphenol A; BPF, bisphenol F; BPS, bisphenol S; IQR, interquartile range.