

Supplemental Table 4. Change From Baseline to Week 26 in Scores for Other Efficacy Endpoints (observed cases)

Endpoint (Full Scale)	Placebo		Lutikizumab		LS Mean Difference (95% CI)	<i>P</i> Value Lutikizumab vs Placebo*
	n	LS Mean (95% CI)	n	LS Mean (95% CI)		
		Lutikizumab vs Placebo*		Lutikizumab vs Placebo*		
AUSCAN stiffness NRS-11, 0–10)	60	−1.9 (−3.0, −0.8)	50	−2.7 (−3.7, −1.6)	−0.8 (−1.6, 0.1)	0.07
Subject assessment of pain intensity, index hand (NRS-11, 0–10)	60	−2.0 (−3.1, −0.9)	50	−2.5 (−3.6, −1.4)	−0.5 (−1.4, 0.4)	0.27
PGA (NRS-11, 0–10)	60	−2.0 (−3.1, −0.9)	50	−2.4 (−3.5, −1.2)	−0.4 (−1.3, 0.6)	0.45
Grip strength, index hand, kg	61	2.1 (−1.6, 5.9)	52	3.0 (−0.9, 6.9)	0.9 (−2.6, 4.4)	0.63
SF-36 (0–100)	60		51			
Physical Component Summary		1.85 (−1.10, 4.79)		3.49 (0.65, 6.33)	1.7 (−0.7, 4.0)	0.17
Mental Component Summary		−0.17 (−3.55, 3.20)		0.40 (−2.97, 3.77)	0.6 (−2.1, 3.2)	0.67
PROMIS SF20a (0–100)	60	1.6 (−0.7, 3.9)	50	2.6 (0.3, 4.8)	1.0 (−0.8, 2.9)	0.28
MHQ, index hand total score (0–100)	60	14.4 (7.2, 21.5)	51	16.0 (8.9, 23.0)	1.6 (−4.1, 7.3)	0.58
Hand function		14.6 (6.7, 22.4)		14.3 (6.6, 22.0)	−0.3 (−6.5, 6.0)	0.93
Activities of daily living		13.7 (5.0, 22.4)		13.8 (5.4, 22.2)	0.1 (−6.8, 7.0)	0.97
Pain		−18.9 (−28.1,		−23.6 (−32.6, −	−4.7 (−12.0, 2.6)	0.20

		–9.6)	14.5)			
Work performance		10.5 (0.2, 20.9)	15.9 (5.7, 26.0)	5.3 (–3.0, 13.6)	0.21	
Aesthetics		10.8 (2.2, 19.4)	6.2 (–2.3, 14.6)	–4.7 (–11.5, 2.2)	0.18	
Satisfaction with hand function		17.1 (6.8, 27.3)	22.2 (12.2, 32.2)	5.1 (–3.0, 13.2)	0.21	
FIHOA	60	0.3 (–1.7, 2.3)	51	–0.8 (–2.8, 1.2)	–1.1 (–2.7, 0.6)	0.20

ANCOVA=analysis of covariance; AUSCAN=Australian/Canadian Osteoarthritis Hand Index; FIHOA, Functional Index for Hand Osteoarthritis; LS=least squares; MHQ=Michigan Hand Outcomes Questionnaire; NRS=numeric rating scale; PGA=patient global assessment of hand osteoarthritis status; PROMIS SF20a=Patient Reported Outcomes Measurement Information System Physical Function Questionnaire; SF-36=36-item short-form health survey.

**P* values are from an ANCOVA model adjusted for treatment groups and country as factors, and including baseline value as a covariate.