

BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<u>http://bmjopen.bmj.com</u>).

If you have any questions on BMJ Open's open peer review process please email <u>info.bmjopen@bmj.com</u>

BMJ Open

Understanding engagement and non-engagement: A longitudinal qualitative study of participant experiences of an exercise referral scheme

Journal:	BMJ Open
Manuscript ID	bmjopen-2018-024370
Article Type:	Research
Date Submitted by the Author:	24-May-2018
Complete List of Authors:	Hanson, Coral; Edinburgh Napier University, School of Health and Social Care; Durham University, Wolfson Research Institute for Health and Wellbeing Physical Activity Special Interest Group Oliver, Emily; Durham University, Department of Sport and Exercise Sciences,; Durham University, Wolfson Research Institute for Health and Wellbeing Physical Activity Special Interest Group Dodd-Reynolds, Caroline; Durham University, Department of Sport and Exercise Sciences; Durham University, Wolfson Research Institute for Health and Wellbeing Physical Activity Special Interest Group Allin, Linda; Northumbria University, Department of Sport, Exercise and Rehabilitation, Faculty of Health and Life Sciences
Keywords:	PUBLIC HEALTH, QUALITATIVE RESEARCH, Health policy < HEALTH SERVICES ADMINISTRATION & MANAGEMENT, PRIMARY CARE, PHYSICAL ACTIVITY, REHABILITATION MEDICINE

SCHOLARONE[™] Manuscripts

For peer review only - http://bmjopen.bmj.com/site/about/guidelines.xhtml

BMJ Open

Understanding engagement and non-engagement: A longitudinal qualitative study of participant experiences of an exercise referral scheme

Coral L Hanson^{1, 2}, (corresponding author), Emily J Oliver^{2, 3}, Caroline J Dodd-Reynolds^{2, 3}, Linda J Allin⁴

¹ School of Health and Social Care, Edinburgh Napier University, Sighthill Campus, Edinburgh, EH11 4DN, UK. Email: c.hanson@napier.ac.uk Tel: 0131 455 3457

² Wolfson Research Institute for Health and Wellbeing Physical Activity Special Interest Group, Durham University, Durham, UK.

³ Department of Sport and Exercise Sciences, Durham University, Durham, UK.

⁴ Department of Sport, Exercise and Rehabilitation, Faculty of Health and Life Sciences, University of Northumbria, Newcastle-upon-Tyne, UK

Word count: 3991

Key Words: Public Health, Qualitative Research, Health Policy, Primary Care, Rehabilitation Medicine, Physical Activity

ABSTRACT

Objectives: Exercise referral schemes are internationally widespread. This study examined experiences of patients referred by healthcare professionals to one such scheme, in order to understand whom it served well, or poorly, and why.

Design: The study employed a qualitative longitudinal approach using semi-structured interviews, with results reported using COREQ guidelines.

Setting: Two leisure centres providing an 'emerging best-practice' exercise referral scheme in northeast England.

Participants: Referred patients (n = 11) who had not yet commenced the scheme, were recruited on a voluntary basis. Seven females and four males, with a range of non-communicable diseases: cardiovascular disease, mental health issues, diabetes, overweight/obesity and musculoskeletal problems participated.

Intervention: 24-week, twice weekly supervised exercise sessions and three one-to-one assessments (pre-scheme, 12-weeks and 24-weeks) for patients referred from primary and secondary care.

Primary outcome measures: Two longitudinal semi-structured interviews, prior to commencement and 12-20 weeks later, were thematically analysed using the framework approach. Analysis comprised seven stages: transcription, familiarisation, coding,

For peer review only - http://bmjopen.bmj.com/site/about/guidelines.xhtml

BMJ Open

development and application of an analytical framework, charting data using a matrix, and
interpretation of data. Interpretation went beyond descriptions of individual cases to develop
themes, which identified and offered possible explanations for differing participant
experiences.

Results: Three overarching themes emerged. First, 'success', with engaged participants focused on health outcomes and reported increases in physical activity. Second, 'struggle', with short-term success but concerns regarding continued engagement. Participants reported scheme dependency and cyclical needs. Finally, 'defeat', where ill health, social anxiety, and/or poor participation experience made engagement difficult.

Conclusion: Some success in engaging those with non-communicable diseases was reported, resulting in positive effects on health and wellbeing. The study highlights complexity within exercise referral schemes, and inequality of access for those with challenging health and social circumstances. Improved, or different, behaviour change support is required for referrals finding engagement difficult.

ARTICLE SUMMARY

Strengths and limitations of this study

- Advancing the predominantly quantitative or cross-sectional literature on participant adherence to exercise referral schemes (ERS), this study explored in-depth experiences over time.
- The study contributes to reducing the evidence gap identified, for example, by the UK National Institute of Health and Care Excellence, about which sub-groups ERSs work for and why.
- Results highlighted the need for discussions about suitability of referrals and delivery practices of this type of intervention, which is internationally widespread.
- The study failed to engage some of the original participants in second interviews, meaning that the experiences of some who may have been the least well-served by the intervention are unknown.
- Results are based on a sample of participants recruited from only one, albeit largescale, ERS and since provider practices may vary, other scheme structures may impact experiences in different ways.

BMJ Open

INTRODUCTION

Regular physical activity (PA) has a beneficial effect on cardiovascular disease risk, diabetes, some cancers and all-cause mortality.¹ Despite this global levels of PA are low, hence the cost of PA to health-care systems in 2013 was estimated to be 53.8 billion international dollars.² Increasing population PA levels is therefore a high priority to reduce non-communicable diseases.³ Understanding participation is important in planning action and this has been widely described in terms of demography, with inequalities apparent.⁴ For example, there is an inverse relationship between PA and indicators of disadvantage such as socio-economic status⁵ and multiple co-morbidities.⁶ PA promotion initiatives must therefore consider how to target the least active.

An understanding of whether current PA programmes disproportionally benefit disadvantaged groups is required. Emerging evidence indicates the converse; with lower socioeconomic status, and increasing number of health conditions, medications and depressive symptoms negatively predicting adherence.⁷ Factors affecting participation are complex, however, with personal and social factors also known to be barriers or facilitators to being active.⁸⁻¹². Understanding how and why existing programmes engage, or do not engage, participants with differing personal circumstance can inform future equitable practice.

An exercise referral scheme (ERS) is one option for health professionals to promote PA for those with non-communicable diseases.¹³ Such schemes are internationally widespread, existing for example, in the United Kingdom (UK),¹⁴ Denmark,¹⁵ Spain¹⁶ and Mexico.¹⁷ In the UK, leisure providers usually deliver ERSs, directing participants into 10-24 weeks of supervised PA. The present study focused on one large scale ERS identified as emerging best

practice by Public Health England.¹⁸ Although ERSs are broadly aimed at those with noncommunicable diseases, there is limited understanding of effective targeting. Exploring whether sub-groups of participants are more or less likely to engage therefore has value in informing practice.¹⁹ This is important because cost-effectiveness analyses indicate that ERSs need to reduce costs by 60%.²⁰ However, lack of evidence about effectiveness for participant sub-groups may have resulted in an underestimation of benefits. Indeed, the UK National Institute for Health and Care Excellence has identified a requirement to understand better what ERS elements work best and for whom.²¹

To implement successful and equitable ERSs, there is a need to better understand who existing programmes work or do not work for, and why. Demographic evidence contributes to understanding the first two of these questions; ^{20 22} however, the third requires a more indepth analytical approach. This study, a longitudinal examination of participant experiences of taking part in an ERS, therefore aims to gain in-depth understanding of differential engagement.

METHODS

The study employed a qualitative approach and longitudinal design to explore experiences of participants referred to a northeast England ERS. Results were reported using the COREQ guidelines.²³ Overarching themes, encompassing a variety of participants with differing social contexts and motives for referral, were established. Northumbria University Faculty of Health and Life Sciences Ethics Committee granted study ethical approval (Ref: 15-03-131781). Participants gave written informed consent.

Context

The ERS received primary and secondary care referrals for those with cardiovascular disease, overweight/obesity, mental health issues, metabolic disease, and musculoskeletal, respiratory and neurological conditions. Previous analysis (n=2233) reported a significant increase in self-reported PA for those who adhered, with being aged \geq 55 years a predictor of successful engagement.²⁴ Scheme design was based on the Transtheoretical Model,^{25 26} (Figure 1). Staff held an industry standard exercise referral qualification.

Figure 1: Scheme Process

The study took place in two of nine leisure centres providing the ERS. Referrals to these leisure centres were representative of the demographic spectrum of participants. This included a broad adult age range, males and females, and a range of economic circumstances and medical conditions. All those referred to the two leisure centres during May and June 2013 (n=102) were eligible to take part.

Patient and Public Involvement

Previous quantitative ERS performance analysis informed the study. Participants were not involved in study design. A results summary was available for study participants.

Sample

Early sampling of participants was by convenience.²⁷ All those invited to attend initial consultations during the first two weeks of the recruitment period (n=25) received an invitation to participate. During initial telephone contact, ERS staff informed referrals that the study consisted of two semi-structured interviews about their ERS experience. The first was conducted prior to starting, the second later in the 24-week period. Postal information was sent to interested referrals, who signed and returned the consent form to register for the study. ERS staff arranged interviews and the researcher had no access to personal details until consent was given. Participants were informed that the researcher was an employee of the scheme provider and that a research objective was to improve service delivery. There was no obligation to take part and ERS involvement was not dependent on this decision. Eight of those initially invited agreed to participate. Later sampling was purposeful, based on emergent themes from earlier initial interviews.²⁸ Recruitment continued until the emergence of new concepts from initial interview analysis ceased.

Data collection and analysis

Data were collected via two longitudinal semi-structured interviews conducted between May and December 2013, which were audio-recorded and transcribed verbatim. CLH, a PhD student who was employed by the scheme provider as a strategic manager, conducted all interviews. Prior to undertaking interviews, CLH completed qualitative interviewing training and received mentoring from LJA, an experienced qualitative researcher. Initial interviews took place in private immediately prior to initial consultations at the participants' leisure venue. One pilot interview using a semi-structured guide was conducted and analysed by CLH and LJA. The guide remained unchanged and the pilot interview deemed suitable for study inclusion. Initial interviews focused on circumstances leading to referral and

Lien

BMJ Open

perceptions of the ERS. Second interviews, which took place 12-20 weeks later, focused on participation or non-participation experiences. Participants checked attendance at sessions with scheme staff and reported this during second interviews. Individual interviews ranged in length from 22-62 minutes (median 48 minutes). Participants checked transcripts for accuracy. Detailed field notes focused on participants' social context, the quality of the interaction and potential researcher bias due to insider knowledge.

Interviews were subject to thematic analysis using the framework approach.²⁹ The use of pseudonyms ensured anonymity. CLH and LJA familiarised themselves with transcripts through reading and rereading, and by listening to audio-recordings to check accuracy. Using manual processes and Microsoft Excel to organise data, they openly recorded preliminary concepts and patterns for three transcripts. After discussion between all authors (n=4), the establishment of agreed codes formed an initial analytical framework. Three more transcripts were analysed before refinement and finalising of the framework to allow comparison within and across all cases. The creation of a matrix allowed for the mapping and exploration of connections within and between participants and categories. During interpretation, analysis went beyond descriptions of individual cases to develop themes identifying and offering possible explanations for types of ERS experience. Participants did not provide feedback on findings, but themes were checked with ERS staff via a workshop.

RESULTS

Participant characteristics

Fifteen referrals took part in initial interviews and 11 completed both interviews. Only participants who completed both interviews were included in the final analysis (Table 1). Four participants did not complete the second interview. Two participants responded stating that they had dropped out and did not have time for the interview due to a new job or caring commitments. The other two did not respond.

Table 1: Participant Characteristics

Participant Pseudonym	Age Group (years)	Gender	Primary reason for referral	Source of referral	Participation status*	Overarching theme
Alice	70+	Female	CVD Secondary Prevention	Cardiac Rehabilitation	Adherer	
Amy	20-29	Female	Overweight/obesity	Primary care	Adherer	Success
Julie	50-59	Female	Overweight/obesity	Primary care	Adherer	Success
Patricia	60-69	Female	Overweight/obesity	Primary care	Adherer	
Brian	60-69	Male	CVD Secondary Prevention	Cardiac Rehabilitation	Adherer	
Margaret	60-69	Female	Mental Health	Primary care	Dropout	Struggle
Peter	50-59	Male	CVD Secondary Prevention	Cardiac Rehabilitation	Adherer	
Paul	50-59	Male	Overweight/obesity	Primary care	Non- starter	
Jackie	40-49	Female	Overweight/obesity	Primary care	Non- starter	
Dorothy	60-69	Female	Musculoskeletal	Primary care	Dropout	Defeat
Dan	50-59	Male	Mental Health	Primary care	Dropout	

*Non-starter: attended initial consultation, but no exercise sessions Dropout: attended initial consultation and some exercise sessions but informed researcher they had stopped attending

Adherer: attended initial consultation and informed researcher they were still attending exercise sessions

Overarching themes

Three overarching themes emerged, each conveying a different referral experience (Figure 2).

Figure 2: Thematic analysis of ERS experiences

For peer review only - http://bmjopen.bmj.com/site/about/guidelines.xhtml

BMJ Open

The first was *success*, with motivated participants focused on health outcomes. The second described a *struggle*, with some level of short-term success. Cyclical changes in circumstances such as health status, and overall concerns regarding continued engagement were evident. The final theme centred on *defeat*, where ill health, social anxiety and/or poor experiences of participation made engagement difficult or unsuccessful. Short excerpts from transcripts give an indication of typical experience, with [...] signifying the joining of different sections. A participant case study illustrates each theme in more depth.

Success: increased physical activity and improved health

Success illustrated how the ERS worked very well for some, with sub-themes of improved health, increased PA and support. These participants tended to have had positive early experiences of sport and were motivated to improve or maintain health. Participation was mainly enjoyable, with peer and/or staff support being important attendance facilitators. Julie highlighted the 'very helpful' staff and how 'enjoying Pilates has motivated me to be coming more'. Although personal goals, for example weight loss, were not always as anticipated, the experience was rewarding and there was a celebration of success. There was an expectation that activity would continue via signposted exit route sessions or independent exercise: 'I will just come on my own. They (other participants) have finished but they still come at the same time' (Patricia).

Illustrating this theme is Alice. She had completed cardiac rehabilitation prior to starting:

Alice: 'I loved sport, I used to cycle to work, then I started doing yoga, but I also like aqua fit, and I love walking around. I couldn't believe when I had a heart attack. [...] You have this fear, I don't walk anywhere where I'm going to fall down and nobody's going to see us. [...] I feel confident about the scheme. I'll move straight on (from cardiac rehabilitation). [...] I'm overweight; I've got to lose at least a stone. [...] You've got to use it or lose it, I've always believed that.'

During her second interview, Alice reported 91% attendance and was very positive. Staff and social support were important in encouraging adherence:

Alice: 'I've really enjoyed it, I feel much healthier again, and I've made loads of new friends. I think the most important thing is I feel that I have got my confidence back. It has definitely lifted all that was frightening. [...] I haven't lost weight but I haven't put any on. I am more content with my life again. More realistic. [...] I love the class, and I like the talking. Yes the mouth exercises, they are very good. [...] It makes you feel as though you belong in a club. [...] (Staff member) is full of fun as well. She does push you along.'

Within the theme of success, there were elements of struggle. Before starting the scheme, there were concerns about how perceived physical limitations and personal situations would affect attendance. Amy discussed self-esteem issues and how she made *'sure that I have got someone with us because I don't feel very confident going out by myself.'*

Participants described using social comparisons³⁰ to make judgments about their personal situation. Both upward and downward comparisons positively reinforced participation. For example, Amy developed positive views of older people's fitness, which encouraged her to

For peer review only - http://bmjopen.bmj.com/site/about/guidelines.xhtml

BMJ Open

do more: 'I've seen what they have got their treadmills on and I'm thinking, I'm only lower than them, I'd best turn it up'. Overall there was a steady improvement in perceptions about ability to be active, and the associated health and social benefits. Enjoyment was both an important facilitator of success and a positive outcome of participation.

Struggle: cyclical needs and scheme dependency

Struggle illustrated how the ERS worked in the short-term for some but highlighted different approaches, or additional measures, may be necessary to encourage sustained increases in PA. Sub-themes of cyclical needs, scheme dependency and multiple barriers indicated that this theme was more complex than *success*. Resulting experiences were more divergent, with difficult life circumstances and/or complex health conditions influencing participation. For these participants, frequently life events had caused a breakdown of their social order (e.g., the death of a loved one or loss of a job), and the ERS enabled a regaining of structure and control. Strugglers perceived the scheme as a way to get lives '*back on track*' (Margaret). This, when combined with complex health problems, meant disengagement could be difficult. Brian, a widower with depression and a history of myocardial infarction, had: '*trouble with my left foot, I am partially blind now, diabetic.* ' He reported a lack of confidence to move on, indicating scheme dependency: '*I'm letting (staff member) set my programme. I might jigger myself up. The scheme is fine. It's ideal. I've been asking him can I stop in it?*' Margaret, in contrast, felt her mental health had improved so she no longer needed to attend but recognised that this was cyclical, and that she may need future support.

Within *struggle*, participants reported increased PA and health gains *'I feel 100% better'* (Brian); however, longer-term positive continuation post-scheme appeared unlikely.

For peer review only - http://bmjopen.bmj.com/site/about/guidelines.xhtml

Peter's experiences illustrate the theme of struggle. For him, social circumstance was particularly influential. His attendance was sporadic (63% of potential sessions). He was unemployed at his first interview and saw a potential return to work as an adherence barrier:

Peter: Well I don't know exactly what is going to happen. I'll just see what it is and how it goes. If I enjoy it, I will stick with it as long as I can. [...] If I find work, I would stop attending.'

During his second interview, it was difficult to gauge his enthusiasm due to his natural reticence: '*I come because it is there and it's available. Or I would be just moping around the house all day doing nothing.*' His intermittent attendance was due to a work-related course and a short period of employment. Cost was an issue:

Peter: 'It's pretty hard because I have got no wages coming in. I've just got Job Seekers Allowance and that doesn't even pay my mortgage, so money is really tight. I did have a little insurance but that money has now run out and I really am starting to struggle. It is fairly cheap, but saying that, when you haven't got a lot of money coming in then it is a lot of money to pay out.'

Peter stated he would continue to attend until he found employment, when he *'might try and come with my partner and play badminton once or twice a week.* ' His enthusiasm for this appeared to be lacking and therefore long-term change in PA was unlikely.

Defeat: inappropriate referral or poor participation experience

For peer review only - http://bmjopen.bmj.com/site/about/guidelines.xhtml

BMJ Open

The third theme was *defeat*. Within this, subthemes of poor health, social anxiety and poor participation experience were apparent. Some participants never attended an exercise session, prevented by ill health (Paul and Dan) or social anxieties: *'The thought of coming here on my own, with nobody else, I like staying in my comfort zone'* (Jackie). Poor participation experience (Dorothy) indicated weaknesses in scheme delivery. Defeated by the barriers faced, they felt ostracised from participation. There was a sense of failure and, for some, shame. For these participants the ERS did not work.

Illustrating this theme is Dorothy, who had been previously referred to the scheme on two occasions. The first time she completed the scheme but did not continue via signposted exit route activities because her friend stopped attending. On the second occasion, she dropped out due to a foot problem. She did not like PA:

Dorothy: 'At school I wasn't very good at PE. I never liked it very much, only did what I had to do. [...] I've been to be the scheme twice before; other than that I don't think I really did any exercise. [...] I've got a real problem with my back. I'm sort of hoping that if I do exercise it will strengthen the muscles in my back and I will be able to do more things.'

During her second interview, Dorothy described how she felt unable to cope with the sessions due to back pain:

Dorothy: 'I did tell her that I'd got a back problem and I was waiting for these injections, but she said 'well start'. [...] I was quite disappointed because I couldn't do much of what they

was asking us to do. It was a class and it was where you do sort of aerobics first and then go to all these sort of stations. I found it really hard.'

Discussing one session, she described a lack of staff support: 'She just said do what you can do and if you can't do whatever it is, just keep your feet moving.' This contrasted with her experience with another staff member: 'She knew when you couldn't do it and she would give you an alternative. So she was really good but she was only there once.'

The scheme had a system of telephone support, but in Dorothy's case, implementation appeared to be lacking:

Dorothy: 'I 'phoned in several times to explain. I left messages but nobody got back to me. I think if maybe someone had 'phoned me back and said 'well come in and you can do the things a different way' it might have encouraged me to go back in again.'

Dorothy raised delivery issues and highlighted the need for a better understanding of protocol implementation by staff. She was very upset by her experience, stating 'It makes you feel like a failure sometimes. I don't think I could go to the doctors and say 'I failed last time can you refer me again'?' Her experience was complex however. She felt unable to access the peer support described by those who engaged successfully. During her first interview, Dorothy described how she found socialising difficult: 'I'm not really a good mixer so I find it quite hard.' During her second interview, she reported that the group felt unwelcoming: 'She was really nice, one lady that was there. Apart from her, I don't think any of the others were welcoming or said anything. Like I said they got into their little twos or threes or whatever.'

BMJ Open

Lack of social confidence contributed to dropout, although in Dorothy's case it does not appear to be have been the primary influence. Defeat illustrates how some participants may struggle to access the peer support identified by others as an important facilitator for adherence.

DISCUSSION

The purpose of this study was to understand experiences of an ERS to give insight regarding what worked, or did not work, to encourage engagement, and for whom. This is important because existing literature questions ERS effectiveness,^{20 22} without exploring adequately how to focus implementation to better personalise support. Three overarching themes emerged. First, *success*, with engaged participants focused on health outcomes. Second, struggle, short-term success but with concerns regarding continued engagement. Finally, *defeat*, where illness, social anxiety, and/or poor participation experience prevented engagement. Within the identified themes, similarities in factors affecting engagement and non-engagement were evident. For those who experienced some measure of success, there were shared enablers such as peer support, achievable and enjoyable activities, staff knowledge and staff/peer support. These are reflected to some extent in systematic review findings of engagement facilitators for ERS.³¹ What this paper adds is insight regarding whom these facilitators worked best for within the ERS context. Specifically, participants who were able to access social support, had positive previous experiences of PA, and were motivated by improving or maintaining their health. Participants often described success in terms of improvements in mental health and self-esteem, perceptions about ability to be active, and the social benefits of participation. Social and psychological benefits were perceived to be as meaningful as measurable physical health benefits, similar to other

reported findings.³² Success illustrated the value of exercise referral for some participants with non-communicable diseases. This was not universal however. The study highlighted unequal abilities to access the scheme, along with differing support requirements, which suggests the need to provide more tailored support for some. The issues identified are reflective of other studies examining barriers to PA irrespective of the presence of a medical condition.⁷¹¹

Adults with complex lives embarked on the ERS with expectations of positive changes in health. While ERS delivery training courses include elements of behaviour change, the training does not appear to be sufficient preparation for staff to deal with identified complex psychological barriers. Indeed the high levels of responsibilities that fitness professionals undertake has led to concerns about adequacy of education and training.³³ Further development of behaviour change elements within national occupational standards for promoting PA could partially address matters. This represents only part of the problem however. This study illustrates how a 'one size fits all' model does not adequately cater for the complex range of referrals received. Indeed, the existing model of universal referral to a common programme is potentially setting such schemes up to fail. This is because current measures of success are typically quantified as uptake and adherence,³⁴⁻³⁶ and/or self-reported changes in PA.^{15 37} Regardless of suitability, providers may feel obliged to 'shoehorn' referrals into schemes if the continuation of funding is reliant on achievement of such key performance indicators. This type of approach fails to consider the complex health and social circumstances of ERS participants, leading to an inadequate focus on what works, and for whom.

BMJ Open

In the case of ERS, there is a need to understand what different approaches are required to support change for those experiencing struggle or defeat. The themes presented in this study may resonate with ERS commissioners and providers and should encourage reflection of approaches to support. *Success* can reinforce good practice, while highlighting potential improvements. *Struggle* can initiate conversations about alternative delivery for those who require more or different support in order to make sustained behaviour change. This may include mechanisms for cyclical support to reengage those who relapse into inactivity and 'weaning' to reduce ERS dependency. Finally, *defeat* can initiate conversations about alternative models of care. At a broad level, approaches may include support from multiple agencies,³⁸ the use of technology,³⁹ or broader system change.⁴⁰

Readers can make choices about whether the identified themes resonate with their own intuitive understanding of such situations, which arguably can improve practice through the process of naturalistic generalization.⁴¹ That said, it is not known whether the experiences of those who declined to participate or dropped out of the study were different to those who took part. For example, we previously established that those under 55 years of age were less likely to engage in the scheme in the first instance and more likely to dropout when they did.²⁴ However, only one participant from this demographic completed a second interview. Additionally, this piece of work did not examine barriers to scheme access for non-starters. Understanding this group, however, is critical for determining who current services are failing and why.

Qualitative analysis is inherently subjective since it is influenced by the assumptions, beliefs and biases of the researcher.⁴² In this case, the researcher was experienced in the management

and delivery of the ERS studied. Potential biases were explored by the use of reflective field notes and in group discussions with all authors. Particular attention was paid to how existing knowledge may have affected discussion with participants and interpretation of results. That said, while in the past an outsider, objective stance was considered desirable in research terms to guard against identification, insider insight can now be considered legitimate and desirable due to the potential for increased empathy with participants.⁴³ After reflection, it was felt that researcher knowledge contributed to the interpretation of data.

CONCLUSION

Overall, the data support arguments that ERSs can disproportionally engage with, and benefit, some disadvantaged groups. Importantly they can successfully engage those with non-communicable diseases, and positively affect health and wellbeing. The value of current ERSs appears to be for those with social confidence and previous positive experiences of PA. Conversely, such schemes may fail those who struggle to access social support due to varying health condition demands, or complex or impaired social circumstances. For those who are unable to adhere, feelings of ostracism and failure may further exacerbate outcome differentials. Ultimately, even programmes that target disadvantaged sub-groups (in the case of ERSs, those with non-communicable diseases) appear at risk of reinforcing inequalities. This study therefore highlights a need for services and systems that better provide for those with dynamic health and social circumstances.

Acknowledgements: Many thanks to ERS staff at Blyth Valley Arts and Leisure for helping with recruitment and organisation of interviews.

BMJ Open

Author Contributions: CLH contributed to the study design, data collection, data analysis and preparation of the final document. EO contributed to data analysis and preparation of the final document. CDR contributed to study design, data analysis and preparation of the final document. LJA contributed to study design, data analysis and preparation of the final document. All authors contributed to this paper and approved the final version.

Funding: Blyth Valley Arts and Leisure was one of two providers of the exercise referral scheme studied and provided funding for a PhD to evaluate the service.

Competing interests: CLH is a former employee of Blyth Valley Arts and Leisure and completed a PhD that was funded by the aforementioned company.

Participant consent: obtained

Ethics approval: Northumbria University Faculty of Health and Life Sciences Ethics Committee (Ref: 15-03-131781)

Provenance and peer review: Not commissioned; externally peer reviewed.

Data sharing statement: No additional data are available.

REFERENCES

 Lee IM, Shiroma EJ, Lobelo F, et al. Effect of physical inactivity on major noncommunicable diseases worldwide: an analysis of burden of disease and life expectancy. *Lancet* 2012;380:219-29. doi: 10.1016/S0140-6736(12)61031-9

2. Ding D, Lawson KD, Kolbe-Alexander TL, et al. The economic burden of physical inactivity: a global analysis of major non-communicable diseases. *Lancet* 2016;388:1311-24. doi: 10.1016/S0140-6736(16)30383-X

 World Health Organisation. *Global action plan for the prevention and control of non-communicable diseases 2013-2020*. Geneva: World Health Organisation, 2013. http://africahealthforum.afro.who.int/IMG/pdf/global_action_plan_for_the_prevention_and_c ontrol_of_ncds_2013-2020.pdf: (Accessed: 05 March 2018)

4. World Health Organisation. *Global status report on noncommunicable diseases*, 2014.
Geneva: World Health Organisation, 2014.
http://apps.who.int/iris/bitstream/10665/148114/1/9789241564854_eng.pdf (Accessed: 05)

March 2018).

5. Eime RM, Charity MJ, Harvey JT, et al. Participation in sport and physical activity: associations with socio-economic status and geographical remoteness. *BMC Public Health* 2015;15:434. doi: 10.1186/s12889-015-1796-0

6. Sievi NA, Senn O, Brack T, et al. Impact of comorbidities on physical activity in COPD. *Respirology* 2015;20(3):413-18. doi: 10.1111/resp.12456

7. Picorelli AMA, Pereira LSM, Pereira DS, et al. Adherence to exercise programs for older people is influenced by program characteristics and personal factors: a systematic review. *Journal of Physiotherapy* 2014;60(3):151-56. doi: 10.1037/0022-006X.51.3.390

BMJ Open

8. Allender S, Cowburn G, Foster C. Understanding participation in sport and physical
activity among children and adults: a review of qualitative studies. Health Educ Res
2006;21(6):826-35. doi: 10.1093/her/cyl063
9. Borodulin K, Sipila N, Rahkonen O, et al. Socio-demographic and behavioral variation in
barriers to leisure-time physical activity. Scand J Public Health 2016;44(1):62-69. doi:
10.1177/1403494815604080
10. Kern ML, Reynolds CA, Friedman HS. Predictors of Physical Activity Patterns Across
Adulthood: A Growth Curve Analysis. Pers Soc Psychol Bulletin 2010. doi:
10.1177/0146167210374834
11. Pan S, Cameron C, DesMeules M, et al. Individual, social, environmental, and physical
environmental correlates with physical activity among Canadians: a cross-sectional study.
BMC Public Health 2009;9(1):21. doi: 10.1186/1471-2458-9-21
12. Zunft H-JF, Friebe D, Seppelt B, et al. Perceived benefits and barriers to physical activity
in a nationally representative sample in the European Union. Public Health Nutr
1999;2(Supplement 1a):153-60. doi: 10.1017/S1368980099000208
13. Sowden S, Raine R. Running along parallel lines: how political reality impedes the
evaluation of public health interventions. A case study of exercise referral schemes in
England. J Epidemiol Community Health 2008;62(9):835 - 41. doi:10.1136/jech.2007.069781
14. Murphy SM, Edwards RT, Williams N, et al. An evaluation of the effectiveness and cost
effectiveness of the National Exercise Referral Scheme in Wales, UK: a randomised
controlled trial of a public health policy initiative. J Epidemiol Community Health
2012;66(8):745-53. doi: 10.1136/jech-2011-200689

15. Sorenson J, Kragstrup J, Stovgaard T, et al. Exercise on Prescription: a randomized study on the effect of counseling vs counseling and supervised exercise. *Scand J Med Sci Sports* 2008;18(3). doi: 10.1111/j.1600-0838.2008.00811.x

16. Gusi N, Reyes M, Gonzalez-Guerrero J, et al. Cost-utility of a walking programme for moderately depressed, obese, or overweight elderly women in primary care: a randomised controlled trial. *BMC Public Health* 2008;8(1):231. doi: 10.1186/1471-2458-8-231

17. Gallegos-Carrillo K, García-Peña C, Salmerón J, et al. Brief Counseling and Exercise Referral Scheme: A Pragmatic Trial in Mexico. *Am J Prev Med* 2017;52(2):249-59.
doi:10.1016/j.amepre.2016.10.021

Public Health England. *Identifying what works for local physical inactivity interventions*.
 London, Public Health England, 2014.

https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/374560/What worksv1 2.pdf: (Accessed: 05 March 2018)

19. Oliver EJ, Hanson CL, Lindsey IA, et al. Exercise on referral: evidence and complexity at the nexus of public health and sport policy. *International Journal of Sport Policy and Politics* 2016:1-6. doi: 10.1186/1471-2458-9-21

20. Campbell F, Holmes M, Everson-Hock E, et al. A systematic review and economic evaluation of exercise referral schemes in primary care: a short report. *Health Technol Assess* 2015;19(60):1-87. doi: 10.3310/hta19600

21. National Institute for Health and Care Excellence. *Physical activity; exercise referral schemes*. National Institute for Health and Care Excellence, 2014. https://www.nice.org.uk/guidance/ph54/resources/physical-activity-exercise-referral-schemes-pdf-1996418406085 (Accessed: 05 March2018).

25 of 32	BMJ Open
	22. Pavey TG, Taylor A, Fox K, et al. Effect of exercise referral schemes in primary care on physical activity and improving health outcomes: systematic review and meta-analysis. <i>BMJ</i>
	2011;343. doi: 10.1136/bmj.d646223. Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research
	(COREQ): a 32-item checklist for interviews and focus groups. <i>Int J Qual Health Care</i> 2007;19(6):349-57. doi: 10.1093/intqhc/mzm042
	24. Hanson CL, Allin LJ, Ellis JG, et al. An evaluation of the efficacy of the exercise on referral scheme in Northumberland, UK: association with physical activity and predictors of
	engagement. A naturalistic observation study. <i>BMJ Open</i> 2013;3(8). doi: 10.1136/bmjopen-2013-002849
	25. Prochaska JO, Marcus BH. <i>The transtheoretical model: Applications to exercise</i>. In:Dishman RK, ed. Advances in exercise adherence. Champaign, IL, England: Human Kinetics 1994:161-80.
	26. Prochaska JO, DiClemente CC. Stages and processes of self-change of smoking: Toward an integrative model of change. <i>J Consult and Clin Psychol</i> 1983;Vol 51(3):390-95. doi: 10.1037/0022-006X.51.3.390
	27. Marshall MN. Sampling for qualitative research. <i>Fam Pract</i> 1996;13(6):522-26. doi:10.1093/fampra/13.6.522
	28. Strauss AL. <i>Qualitative analysis for social scientists</i> . Cambridge: Cambridge University Press, 1987.
	29. Spencer L, Ritchie J, Lewis J, et al. <i>Quality in qualitative evaluation: a framework for assessing research evidence</i> . National Centre for Social Research, 2003.

http://dera.ioe.ac.uk/21069/2/a-quality-framework-tcm6-38740.pdf (Accessed: 05 March 2018).

30. Festinger L. A Theory of Social Comparison Processes. *Human Relations* 1954;7(2):117-40. doi: 10.1177/001872675400700202

31. Morgan F, Battersby A, Weightman AL, et al. Adherence to exercise referral schemes by participants – what do providers and commissioners need to know? A systematic review of barriers and facilitators. *BMC Public Health* 2016;16(1):1-11 doi: 10.1186/s12889-016-2882-7

32. Pentecost C, Taket A. Understanding exercise uptake and adherence for people with chronic conditions: a new model demonstrating the importance of exercise identity, benefits of attending and support. *Health Educ Res* 2011;26(5):908-22. doi: 10.1093/her/cyr052

33. De Lyon ATC, Neville RD, Armour KM. The Role of Fitness Professionals in Public Health: A Review of the Literature. *Quest* 2017;69(3):313-30. doi:

10.1080/00336297.2016.1224193

34. Dugdill L, Graham R, McNair F. Exercise referral: the public health panacea for physical activity promotion? A critical perspective of exercise referral schemes; their development and evaluation. *Ergonomics* 2005;48:1390-410. doi: 10.1080/00140130500101544

35. Gidlow C, Johnston LH, Crone D, et al. Socio-demographic patterning of referral, uptake and attendance in Physical Activity Referral Schemes. *J Public Health* 2007;29(2):107-13. doi:10.1093/pubmed/fdm002

36. Sowden S, Breeze E, Barber J, et al. Do general practices provide equitable access to physical activity interventions? *Br J Gen Pract* 2008;58:e1-e8. doi: 10.3399/bjgp08X342237


BMJ Open

37. Duda JL, Williams GC, Ntoumanis N, et al. Effects of a standard provision versus an
autonomy supportive exercise referral programme on physical activity, quality of life and
well-being indicators: a cluster randomised controlled trial. Int J Behav Nutr Phys Act
2014;11(1):1-15. doi: 10.1186/1479-5868-11-10
38. Sugai G, Horner RH, Dunlap G, et al. Applying positive behavior support and functional
behavioral assessment in schools. J Posit Behav Interv 2000;2(3):131-43. doi:
10.1177/109830070000200302
39. Gallagher R, Gallagher P, Zhang L, et al. Technology Use and Confidence in Cardiac
Rehabilitation Patients. Heart Lung Circ 2016;25:S333. doi: 10.1016/j.hlc.2016.06.791
40. Michie S, van Stralen MM, West R. The behaviour change wheel: A new method for
characterising and designing behaviour change interventions. Implement Sci 2011;6(1):42.
doi: 10.1186/1748-5908-6-42
41. Stake RE, Trumbull DJ. Naturalistic generalizations. Review Journal of Philosophy and
Social Science 1982;7(1):1-12.

42. Creswell JW, Miller DL. Determining validity in qualitative inquiry. *Theory into practice* 2000;39(3):124-30. doi: 10.1207/s15430421tip3903

43. Gair S. Feeling Their Stories: Contemplating Empathy, Insider/Outsider Positionings, and Enriching Qualitative Research. *Qual Health Res* 2012;22(1):134-43. doi:

10.1177/1049732311420580



60




Figure 1: Scheme Process

266x370mm (300 x 300 DPI)




Figure 2: Thematic analysis of ERS experiences

206x148mm (300 x 300 DPI)

No	Item	Guide questions/description	Information	Reported in manuscript
	Domain 1: Resea	arch team and reflexivity		
	Personal Character			
1	Interviewer/ facilitator:	Which author/s conducted the interview or focus group?	CLH	\checkmark
2	Credentials	What were the researcher's credentials?	CLH PhD student and experienced in working in intervention studied. LJA experienced qualitative researcher	N
3	Occupation	What was their occupation at the time of the study?	PhD student and strategic manager in the ERS provider organisation	N
4	Gender	Was the researcher male or female?	Female (referred to as she)	\checkmark
5	Experience and training	What experience or training did the researcher have?	CLH: Qualitative research training and support from experienced researcher (LJA)	V
	Relationship with			
6	Relationship established	Was a relationship established prior to study commencement?	No, participants were approached by ERS staff to explain the study and seek consent.	V
7	Participant knowledge of the interviewer	What did the participants know about the researcher? e.g. personal goals, reasons for doing the research	Participants were informed of researcher's employment status and that the research aimed to improved service delivery	V
8	Interviewer characteristics	What characteristics were reported about the interviewer/facilitator? <i>e.g.</i> <i>Bias, assumptions, reasons</i> <i>and interests in the research</i> <i>topic</i>	Acknowledgement of potential bias due to insider knowledge. Interest in service improvement and employment status of interviewer disclosed in study invitation information.	V
	Domain 1: Study	y design	·	
	Theoretical frame			
9	Methodological orientation and Theory	What methodological orientation was stated to underpin the study? <i>e.g.</i> grounded theory, discourse analysis, ethnography, phenomenology, content analysis	Thematic analysis using the framework approach	N
	Participant select			ι
10	Sampling	How were participants selected? <i>e.g. purposive,</i> <i>convenience, consecutive,</i>	Initially convenience from a defined group, then purposive	V

COREQ GUIDELINES REPORTING CHECKLIST: Understanding engagement and non-engagement: A longitudinal qualitative study of participant experiences of an exercise referral scheme

1 2
3
4 5
6 7
7 8 9
10
12 13
14 15
16 17
14 15 16 17 18 19
20 21
22 23
24 25
26 27
 19 20 21 22 23 24 25 26 27 28 29 30
29 30 31
32 33
33 34 35
35 36 37
38
39 40
41 42
43 44
45 46
47 48
49 50
51 52
53 54
55 56
57 58
59

		snowball		
11	Method of	How were participants	By telephone by scheme	
	approach	approached? e.g. face-to-face,	provider	
	off the	telephone, mail, email	<u>r</u>	
12	Sample size	How many participants were	15 initially but only 11	
12	Sumple Size	in the study?	completed both interviews	•
13	Non-	How many people refused to	Of initial invitees 13/25	
15	participation	participate or dropped out?	refused. Of 15 initial	v
	participation	Reasons?	participants, 4 dropped out.	
		ixeasons!	(no response n=2, too busy	
			(no response n-2, too busy n=2)	
	Setting		n-2)	
14	Setting of data	Where was the data collected?	Leisure centre where ERS	2
14	collection		was attended	v
1.5		e.g. home, clinic, workplace		
15	Presence of non-	Was anyone else present	No, interviews were	
	participants	besides the participants and	conducted in private	
1.0		researchers?		
16	Description of	What are the important	Gender, age group, medical	\mathcal{N}
	sample	characteristics of the sample?	reason for referral and date	
		e.g. demographic data, date	range for interviews	
			reported	
	Data collection		~ · · · ·	
17	Interview guide	Were questions, prompts,	Semi-structured interview	
		guides provided by the	guide used. Pilot tested	
		authors? Was it pilot tested?		
18	Repeat	Were repeat interviews carried	Yes, one further interview	
	interviews	out? If yes, how many?	after 12-20 weeks	
19	Audio/visual	Did the research use audio or	Yes, the interviews were	
	recording	visual recording to collect the	audio recorded	
		data?		
20	Field notes	Were field notes made during	Yes detailed field notes	
		and/or after the interview or	were made directly after	
		focus group?	interviews	
21	Duration	What was the duration of the	Range and median length	
		interviews or focus groups?	reported: 22-62 minutes	
			(median 48 minutes)	
22	Data saturation	Was data saturation discussed?	Recruitment stopped when	
			no new themes were	
			emerging	
23	Transcripts	Were transcripts returned to	Transcripts were checked	
	returned	participants for comment	by participants.	
		and/or correction?	- 1 1	
	Domain 3: analy		· · ·	
	Data analysis	2		
24	Number of data	How many data coders coded	N=2 (CLH and LJA)	
	coders	the data?	independently	
			N=4 in total at data	
			workshops	
25	Description of	Did authors provide a	Yes figure 2 visually	
-	the coding tree	description of the coding tree?	describes the coding tree	•
26	Derivation of	Were themes identified in	Identified from data	
_0	themes	advance or derived from the		,
		data?		
27	Software	What software, if applicable,	No software used, data	

28 Participant checking Did participants provide feedback on the findings? No, but themes were checked with ERS staff at a workshop. 29 Quotations presented Were participant quotations presented to illustrate the themes / findings? Was each quotation identified? e.g. participant number Yes, participants identified using a pseudonym √ 30 Data and findings consistent Was there consistency between findings? Themes were illustrated by participant quotations √ 31 Clarity of major the data presented in the findings? Were major themes clearly presented in the findings? Three major experience themes were identified success, struggle and defeat √ 32 Clarity of minor themes Is there a description of minor themes? Minor theme: For success: improved health, increased PA, enjoyment and support For struggle: scheme dependency, multiple barriers and cyclical needs For defeat: inappropriate referral, poor participation experience, social anxiety. Diverse cases discussed within themes
29 Quotations presented Were participant quotations presented to illustrate the themes / findings? Was each quotation identified? e.g. participant number Yes, participants identified using a pseudonym 30 Data and findings consistent Was there consistency between the data presented and the findings? Themes were illustrated by participant quotations 31 Clarity of major themes Were major themes clearly presented in the findings? Three major experience themes were identified 32 Clarity of minor themes Is there a description of diverse cases or discussion of minor themes? Minor themes identified within each major theme: For success: improved health, increased PA, enjoyment and support For struggle: scheme dependency, multiple barriers and cyclical needs For defeat: inappropriate referral, poor participation experience, social anxiety. Diverse cases discussed
presented presented to illustrate the themes / findings? Was each quotation identified? e.g. participant number using a pseudonym 30 Data and findings consistent Was there consistency between the data presented and the findings? Themes were illustrated by participant quotations 31 Clarity of major themes clearly presented in the findings? Were major themes clearly presented in the findings? Three major experience themes were identified: success, struggle and defeat 32 Clarity of minor themes? Is there a description of diverse cases or discussion of minor themes? Minor themes identified view in the alth, increased PA, enjoyment and support For struggle: scheme dependency, multiple barriers and cyclical needs For defeat: inappropriate referral, poor participation experience, social anxiety. Diverse cases discussed
findings consistent the data presented and the findings? participant quotations 31 Clarity of major themes Were major themes clearly presented in the findings? Three major experience themes were identified: success, struggle and defeat 32 Clarity of minor themes Is there a description of diverse cases or discussion of minor themes? Minor themes identified within each major theme: For success: improved health, increased PA, enjoyment and support For struggle: scheme dependency, multiple barriers and cyclical needs For defeat: inappropriate referral, poor participation experience, social anxiety. Diverse cases discussed
31 Clarity of major themes Were major themes clearly presented in the findings? Three major experience themes were identified: success, struggle and defeat 32 Clarity of minor themes Is there a description of diverse cases or discussion of minor themes? Minor themes identified within each major theme: For success: improved health, increased PA, enjoyment and support For struggle: scheme dependency, multiple barriers and cyclical needs For defeat: inappropriate referral, poor participation experience, social anxiety. Diverse cases discussed
32 Clarity of minor themes Is there a description of diverse cases or discussion of minor themes? Minor themes identified within each major theme: For success: improved health, increased PA, enjoyment and support For struggle: scheme dependency, multiple barriers and cyclical needs For defeat: inappropriate referral, poor participation experience, social anxiety. Diverse cases discussed

BMJ Open

How do participant experiences and characteristics influence engagement in exercise referral? A qualitative longitudinal study of a scheme in Northumberland, United Kingdom.

Journal:	BMJ Open
Manuscript ID	bmjopen-2018-024370.R1
Article Type:	Research
Date Submitted by the Author:	09-Oct-2018
Complete List of Authors:	Hanson, Coral; Edinburgh Napier University, School of Health and Social Care; Durham University, Wolfson Research Institute for Health and Wellbeing Physical Activity Special Interest Group Oliver, Emily; Durham University, Department of Sport and Exercise Sciences,; Durham University, Wolfson Research Institute for Health and Wellbeing Physical Activity Special Interest Group Dodd-Reynolds, Caroline; Durham University, Department of Sport and Exercise Sciences; Durham University, Wolfson Research Institute for Health and Wellbeing Physical Activity Special Interest Group Allin, Linda; Northumbria University, Department of Sport, Exercise and Rehabilitation, Faculty of Health and Life Sciences
Primary Subject Heading :	Public health
Secondary Subject Heading:	Rehabilitation medicine, Qualitative research
Keywords:	PUBLIC HEALTH, QUALITATIVE RESEARCH, Health policy < HEALTH SERVICES ADMINISTRATION & MANAGEMENT, PRIMARY CARE, PHYSICAL ACTIVITY, REHABILITATION MEDICINE

SCHOLARONE[™] Manuscripts

BMJ Open

1	How do participant experiences and characteristics influence engagement in exercise
2	referral? A qualitative longitudinal study of a scheme in Northumberland, United
3	Kingdom.
4	
5	Coral L Hanson ^{1, 2} , (corresponding author), Emily J Oliver ^{2, 3} , Caroline J Dodd-Reynolds ² ,
6	³ , Linda J Allin ⁴
7	
8	¹ School of Health and Social Care, Edinburgh Napier University, Sighthill Campus,
9	Edinburgh, EH11 4DN, UK. Email: c.hanson@napier.ac.uk Tel: 0131 455 3457
10	
11	² Wolfson Research Institute for Health and Wellbeing Physical Activity Special Interest
12	Group, Durham University, Durham, UK.
13	
14	³ Department of Sport and Exercise Sciences, Durham University, Durham, UK.
15	
16	⁴ Department of Sport, Exercise and Rehabilitation, Faculty of Health and Life Sciences,
17	University of Northumbria, Newcastle-upon-Tyne, UK
18	
19	Word count: 3991
20	
21 22	Key Words: Public Health, Qualitative Research, Health Policy, Primary Care, Rehabilitation Medicine, Physical Activity
22	Medicine, Fnysical Activity
	1 For peer review only - http://bmjopen.bmj.com/site/about/guidelines.xhtml

2	
3	
4	
5	
6	
7	
8	
9	
10	
11	
12	
13	
14	
15	
16	
17	
18	
19	
20	
∠∪ ว1	
21 22	
22	
23	
24	
25	
26	
27	
28	
29	
30	
31	
32	
33	
34	
35	
22	
36 27	
37 38	
39	
40	
41	
42	
43	
44	
45	
46	
47	
48	
49	
50	
51	
52	
52 53	
53 54	
56	
57	
58	
59	
60	

ABSTRACT

2 **Objectives:** Exercise referral schemes are internationally widespread. This study aimed to

- 3 gain an insight into differential engagement through understanding participant experiences of
- 4 patients referred by healthcare professionals to one such scheme in the United Kingdom

5

1

6 **Design:** The study employed a qualitative longitudinal approach using semi-structured

7 interviews, with results reported using COREQ guidelines.

8

9 Setting: Two leisure centres providing an 'emerging best-practice' exercise referral scheme
10 in northeast England.

11

Participants: Referred patients (n = 11) who had not yet commenced the scheme, were
recruited on a voluntary basis. Seven females and four males, with a range of noncommunicable diseases: cardiovascular disease, mental health issues, diabetes,
overweight/obesity and musculoskeletal problems participated.

16

Intervention: 24-week, twice weekly supervised exercise sessions and three one-to-one
assessments (pre-scheme, 12-weeks and 24-weeks) for patients referred from primary and
secondary care.

20

Primary outcome measures: Two longitudinal semi-structured interviews, prior to
commencement and 12-20 weeks later, were thematically analysed using the framework
approach. Analysis comprised seven stages: transcription, familiarisation, coding,

e 3 of 41		BMJ Open
	1	development and application of an analytical framework, charting data using a matrix, and
	2	interpretation of data. Interpretation went beyond descriptions of individual cases to develop
	3	themes, which identified and offered possible explanations for differing participant
	4	experiences.
	5	
	6	Results: Three overarching themes emerged. First, 'success', with engaged participants
	7	focused on health outcomes and reported increases in physical activity. Second, 'struggle',
	8	with short-term success but concerns regarding continued engagement. Participants reported
	9	scheme dependency and cyclical needs. Finally, 'defeat', where ill health, social anxiety,
	10	and/or poor participation experience made engagement difficult.
	11	
	12	Conclusion: Some success in engaging those with non-communicable diseases was reported,
	13	resulting in positive effects on health and wellbeing. The study highlights complexity within
	14	exercise referral schemes, and inequality of access for those with challenging health and
	15	social circumstances. Improved, or different, behaviour change support is required for
	16	referrals finding engagement difficult.

1	ARTICLE SUMMARY
2	Strengths and limitations of this study
3	
4	> This study advances the predominantly quantitative literature on participant adherence
5	to exercise referral by using a longitudinal qualitative design to gain a deeper
6	understanding of the experience of patients with non-communicable diseases referred
7	to an exercise referral scheme (ERS).
8	
9	> The study provides insight into the complexity of ERS engagement and the
10	experiences of a group that has been little researched; those who did not successfully
11	engage with the ERS.
12	
13	> The study was unable to engage some of the original participants in second
14	interviews, meaning that the experiences of some who may have been least well-
15	served by the intervention are unknown.
16	
17	> The sample of participants were recruited from only one, albeit large-scale, ERS,
18	meaning that findings relate to this particular scheme and sample.
19	
20	Qualitative interviews can only provide information on what participants recall or are
21	prepared to reveal about their perceived experiences within a particular interview
22	context, meaning that the potential for recall bias is always present.
23	
24	
25	
	4 For peer review only - http://bmjopen.bmj.com/site/about/guidelines.xhtml

BMJ Open

1 INTRODUCTION

Regular physical activity (PA) has a beneficial effect on cardiovascular disease risk, diabetes, some cancers and all-cause mortality.¹ The global cost of inactivity to health-care in 2013 was estimated to be 53.8 billion international dollars² and therefore increasing PA levels is a high priority to reduce non-communicable diseases.³ Participation in PA has been widely described in terms of demography, with inequalities apparent.⁴ For example, there is an inverse relationship between PA and indicators of disadvantage such as socio-economic status⁵ and multiple co-morbidities.⁶ In order to have the greatest impact, PA promotion initiatives must therefore consider the context, and barriers and facilitators to engagement specifically in disadvantaged populations. Emerging evidence indicates that current PA programmes can fail to engage or retain more

disadvantaged participants. Lower socioeconomic status, and increasing number of health
conditions, medications and depressive symptoms have been reported to negatively predict
adherence.⁷ Factors affecting participation are complex, however, with personal and social
factors such as positive childhood PA experience and social support for PA known to
positively influence activity levels.⁸⁻¹² Understanding how and why existing programmes
engage, or do not engage, participants with differing personal circumstance can inform future
equitable practice.

An exercise referral scheme (ERS) is one option for health professionals to promote PA for
those with non-communicable diseases.¹³ Such schemes are internationally widespread,
existing for example, in the United Kingdom (UK),¹⁴ Denmark,¹⁵ Spain¹⁶ and Mexico.¹⁷ In
the UK, leisure providers usually deliver ERSs, directing participants into 10-24 weeks of

2	
3	
4 5	
6	
7	
8	
9	
10	
11 12	
13 14	
15	
16 17 18	
17 18	
19	
20	
21	
22	
23 24	
24 25	
26 27	
28	
29 30	
30 31	
32	
32 33	
34	
35	
36 37	
38	
39	
40	
41 42	
42 43	
44	
45	
46	
47 48	
48 49	
50	
51	
52	
53 54	
54 55	
56	
57	
58	
59 60	
00	

1

1	supervised PA. The present study focused on one large scale ERS identified as emerging best
2	practice by Public Health England. ¹⁸ Although ERSs are broadly aimed at those with non-
3	communicable diseases, there is limited understanding of effective targeting. Exploring
4	whether sub-groups of participants are more or less likely to engage therefore has value in
5	informing practice. ¹⁹ This is important because cost-effectiveness analyses indicate that ERSs
6	need to reduce costs by 60%. ²⁰ However, lack of evidence about effectiveness for participant
7	sub-groups may have resulted in an underestimation of benefits. Indeed, the UK National
8	Institute for Health and Care Excellence has identified a requirement to understand better
9	what ERS elements work best and for whom. ²¹
10	
11	To implement successful and equitable ERSs, there is a need to better understand who
12	existing programmes work or do not work for. Demographic evidence contributes some
13	knowledge; ^{20 22} but a more in-depth analytical approach is required to increase understanding
14	of other factors influencing engagement. This longitudinal qualitative study aimed to gain an
15	insight into differential engagement through understanding participant experiences of an
16	ERS.
17	
18	METHODS
19	
20	The study employed longitudinal qualitative design to explore experiences of participants
21	referred to the Northumberland ERS. Results were reported using the COREQ guidelines. ²³
22	Overarching themes, encompassing a variety of participants with differing social contexts and
23	motives for referral, were established. Northumbria University Faculty of Health and Life
24	Sciences Ethics Committee granted study ethical approval (Ref: 15-03-131781). Participants
25	gave written informed consent.

BMJ Open

1	
2	Context
3	
4	The ERS received primary and secondary care referrals for those with cardiovascular disease,
5	overweight/obesity, mental health issues, metabolic disease, and musculoskeletal, respiratory
6	and neurological conditions. Previous analysis (n=2233) reported a significant increase in
7	self-reported PA for those who adhered, with being aged \geq 55 years a predictor of successful
8	engagement. ²⁴ Scheme design was based on the Transtheoretical Model. ^{25 26} It consisted of
9	three one-to-one consultations and 24 weeks of twice-weekly PA sessions (Figure 1). During
10	consultations, participants chose which PA sessions to attend. Those who did not attend
11	activity sessions for one week were contacted by telephone or post. Each ERS session cost
12	£3.40. Participants could purchase a discounted direct debit fitness and swimming
13	membership while taking part and after completion (£24.00/month). Staff held an industry
14	standard exercise referral qualification.
15	
16	Figure 1: Scheme Process
17	
18	The study took place in two of nine leisure centres providing the ERS. Referrals to these
19	leisure centres were representative of the demographic spectrum of participants. This
20	included a broad adult age range, males and females, and a range of economic circumstances
21	and medical conditions. All those referred to the two leisure centres during May and June
22	2013 (n=102) were eligible to take part.
23	
24	Patient and Public Involvement
25	
	7 For peer review only - http://bmjopen.bmj.com/site/about/guidelines.xhtml

Previous binary logistic regression analysis of demographic and personal factors associated with engagement and adherence to the ERS^{24} informed the study. Participants were not involved in study design. A results summary was available for study participants. Sample The Northumberland ERS provided a convenient sample,-²⁷ which was easily accessible to CLH, given her employment. All those invited to attend initial consultations during the first two weeks of the recruitment period (n=25) received an invitation to participate. During initial telephone contact, ERS staff informed referrals that the study consisted of two semi-structured interviews about their ERS experience. The first was conducted prior to starting, the second later in the 24-week period. Postal information was sent to interested referrals, who signed and returned the consent form to register for the study. ERS staff arranged interviews and the researcher had no access to personal details until consent was given. Participants were informed that the researcher was an employee of the scheme provider and that a research objective was to improve service delivery. There was no obligation to take part and ERS involvement was not dependent on this decision. Eight of those initially invited agreed to participate. Later sampling was purposeful, based on developing themes (those with multiple medical conditions and referrals under 50 years old) from earlier initial interviews.²⁸ ERS staff were asked to invite referrals with only these characteristics to take part later in the study. Recruitment continued until no new -overarching themes developed from initial interview analysis. Data collection and analysis

Page 9 of 41

1

BMJ Open

1 2	
2	
3 ⊿	
4 5 6	
5	
/ ያ	
a	
7 8 9 10	
11	
12	
13	
11 12 13 14 15 16 17	
15	
16	
17	
17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33	
19	
20	
21	
22	
23	
24	
25	
20 27	
27 28	
20 29	
30	
31	
32	
33	
34 35 36 37 38	
35	
36	
37	
38	
39	
40	
41	
42 43	
43 44	
44 45	
46	
47	
48	
49	
50	
51	
52	
53	
54	
55	
56 57	
57	
58 59	
59 60	
00	

1	Data were collected via two longitudinal semi-structured interviews conducted between May
2	and December 2013, which were audio-recorded and transcribed verbatim. CLH, a PhD
3	student who was employed by the scheme provider as a strategic manager, conducted all
4	interviews. CLH had 15 years' experience of working for the ERS but was not involved in
5	delivery during the study. Prior to undertaking interviews, CLH completed qualitative
6	interviewing training and received mentoring from LJA, an experienced qualitative
7	researcher. Initial interviews took place in private immediately prior to initial consultations at
8	the participants' leisure venue. One pilot interview using a semi-structured guide
9	(supplementary file 1) was conducted and analysed by CLH and LJA. Topics covered
10	included PA history, motivators for referral, perceptions and expectations of the ERS, and
11	perceived barriers and facilitators to taking part. The guide remained unchanged and the pilot
12	interview deemed suitable for study inclusion. Initial interviews focused on circumstances
13	leading to referral and perceptions of the ERS. Second interviews, which took place 12-20
14	weeks later, focused on participation or non-participation experiences (supplementary file 2).
15	Participants checked attendance at sessions with scheme staff and reported this during second
16	interviews. Individual interviews ranged in length from 22-62 minutes (median 48 minutes).
17	Participants checked transcripts for accuracy. Detailed field notes focused on participants'
18	social context, the quality of the interaction and potential researcher bias due to insider
19	knowledge.
20	

20

Interviews were subject to thematic analysis using the framework approach.²⁹ The use of
pseudonyms ensured anonymity. CLH and LJA familiarised themselves with transcripts
through reading and rereading, and by listening to audio-recordings to check accuracy. Using
manual processes and Microsoft Excel to organise data, they openly recorded preliminary
concepts and patterns for three transcripts. After discussion between all authors (n=4), the

1	establishment of agreed codes formed an initial analytical framework. Three more transcripts						
2	were analysed before refinement and finalising of the framework to allow comparison within						
3	and across all cases. The creation of a matrix allowed for the mapping and exploration of						
4	connections within and between participants and categories. During interpretation, analysis						
5	went beyond descriptions of individual cases to develop themes identifying and offering						
6	possible explanations for types of ERS experience. Participants did not provide feedback on						
7	findings, but themes were checked with ERS staff via a workshop.						
8							
9	RESULTS						
10	RESULTS Participant characteristics						
11	Participant characteristics						
12							
13	Fifteen referrals took part in initial interviews and 11 completed both interviews. Only						
14	participants who completed both interviews were included in the final analysis (Table 1).						
15	Four participants did not complete the second interview. Two participants responded stating						
16	that they had dropped out and did not have time for the interview due to a new job or caring						
17	commitments. The other two did not respond. Three of the four were under 50 years old.						
18							
19	Table 1: Participant Characteristics						
	Participant Age Primary reason Source of Self-reported PA Employ- ment Status* theme						

Participant Pseudonym	Age Group (years)	Gender	Primary reason for referral	Source of referral	Self-reported PA history	Employ- ment status	Participation status*	Overarching theme
Alice	70+	Female	CVD Secondary Prevention	Cardiac Rehabilitation	Previously very active, enjoyed PA	Retired	Adherer	
Amy	20-29	Female	Overweight/ obesity	Primary care	Previously active, enjoyed PA	Home- maker	Adherer	Success
Julie	50-59	Female	Overweight/ obesity	Primary care	Previously very active, enjoyed PA	Employed	Adherer	

BMJ Open

Patricia	60-69	Female	Overweight/ obesity	Primary care	Previously active, enjoyed PA	Retired	Adherer	
Brian	60-69	Male	CVD Secondary Prevention	Cardiac Rehabilitation	Previously active, enjoyed PA	Retired	Adherer	
Margaret	60-69	Female	Mental Health	Primary care	Intermittently active, enjoyed PA but hated sport	Retired	Dropout	Struggle
Peter	50-59	Male	CVD Secondary Prevention	Cardiac Rehabilitation	Previously active, enjoyed PA	Un- employed	Adherer	
Paul	50-59	Male	Overweight/ obesity	Primary care	Previously active, enjoyed PA	Receiving disability benefit	Medically excluded	
Jackie	40-49	Female	Overweight/ obesity	Primary care	Previously inactive, disliked PA	Carer	Non- attender	Defeat
Dorothy	60-69	Female	Musculoskeletal	Primary care	Previously inactive, disliked PA	Retired	Dropout	
Dan	50-59	Male	Mental Health	Primary care	Previously active, enjoyed PA	Employed	Dropout (health reason)	

*Medically excluded: attended initial consultation but was excluded from scheme participation due to medical reasons (physiological measures above scheme acceptance guidelines e.g. blood pressure \geq 180/100 mmHg or resting heart rate \geq 100 beats per minute) Non-attender: attended initial consultation, but no exercise sessions

2.04

Dropout: attended initial consultation and some exercise sessions but informed researcher they had stopped attending **Adherer:** attended initial consultation and informed researcher they were still attending exercise sessions

Overarching themes

5 Three overarching themes emerged, each conveying a different referral experience (Figure 2).

7 Figure 2: Thematic analysis of ERS experiences

- 9 The first was *success*, with motivated participants focused on health outcomes. The second
- 10 described a *struggle*, with some level of short-term success. Cyclical changes in
- 11 circumstances such as health status, and overall concerns regarding continued engagement
- 12 were evident. The final theme centred on *defeat*, where ill health, social anxiety and/or poor
- 13 experiences of participation made engagement difficult or unsuccessful. Short excerpts from

1	transcripts give an indication of typical experience, with [] signifying the joining of
2	different sections. A participant case study illustrates each theme in more depth.
3	
4	Success: increased physical activity and improved health
5	
6	Success illustrated how the ERS worked very well for some, with sub-themes of improved
7	health, increased PA and support. These participants tended to have had positive early
8	experiences of sport and were motivated to improve or maintain health. Participation was
9	mainly enjoyable, with peer and/or staff support being important attendance facilitators. Julie
10	highlighted the 'very helpful' staff and how 'enjoying Pilates has motivated me to be coming
11	more'. Although personal goals, for example weight loss, were not always as anticipated, the
12	experience was rewarding and there was a celebration of success. There was an expectation
13	that activity would continue via signposted exit route sessions or independent exercise: 'I will
14	just come on my own. They (other participants) have finished but they still come at the same
15	time' (Patricia).
16	
17	Illustrating this theme is Alice. She had completed cardiac rehabilitation prior to starting:
18	
19	Alice: 'I loved sport, I used to cycle to work, then I started doing yoga, but I also like aqua
20	fit, and I love walking around. I couldn't believe when I had a heart attack. [] You have this
21	fear, I don't walk anywhere where I'm going to fall down and nobody's going to see us. [] I
22	feel confident about the scheme. I'll move straight on (from cardiac rehabilitation). [] I'm
23	overweight; I've got to lose at least a stone. [] You've got to use it or lose it, I've always
24	believed that.'
25	
	12

For peer review only - http://bmjopen.bmj.com/site/about/guidelines.xhtml

BMJ Open

2	
3	
4	
5	
6	
7	
8	
9	
10	
11	
12	
13	
14	
15	
16	
16 17	
18	
19	
20	
21	
22	
23	
24	
25	
26	
27	
28	
29	
30	
31	
32	
33	
34	
35	
36	
37	
38	
39	
40	
41	
42	
43	
44	
45	
46	
47	
48	
49	
50	
51	
52	
53	
54	
55	
56	
57	
58	
59	
60	

During her second interview, Alice reported 91% attendance and was very positive. Staff and
 social support were important in encouraging adherence:

3

Alice: 'I've really enjoyed it, I feel much healthier again, and I've made loads of new friends.
I think the most important thing is I feel that I have got my confidence back. It has definitely
lifted all that was frightening. [...] I haven't lost weight but I haven't put any on. I am more
content with my life again. More realistic. [...] I love the class, and I like the talking. Yes the
mouth exercises, they are very good. [...] It makes you feel as though you belong in a club.
[...] (Staff member) is full of fun as well. She does push you along.'

10

Within the theme of success, there were elements of struggle. Before starting the scheme, there were concerns about how perceived physical limitations and personal situations would affect attendance. Amy discussed self-esteem issues and how she made '*sure that I have got someone with us because I don't feel very confident going out by myself.*'

15

Participants described using social comparisons³⁰ to make judgments about their personal
situation. Both upward and downward comparisons positively reinforced participation. For
example, Amy developed positive views of older people's fitness, which encouraged her to
do more: *Tve seen what they have got their treadmills on and I'm thinking, I'm only lower than them, I'd best turn it up'*. Overall there was a steady improvement in perceptions about
ability to be active, and the associated health and social benefits. Enjoyment was both an
important facilitator of success and a positive outcome of participation.

23

25

24 Struggle: cyclical needs and scheme dependency

2		
- 3 4	1	Struggle illustrated how the ERS worked in the short-term for some but highlighted different
5	2	approaches, or additional measures, may be necessary to encourage sustained increases in
7 8	3	PA. Sub-themes of cyclical needs, scheme dependency and multiple barriers indicated that
9 10	4	this theme was more complex than success. Resulting experiences were more divergent, with
11 12	5	difficult life circumstances and/or complex health conditions influencing participation. For
13 14	6	these participants, frequently life events had caused a breakdown of their social order (e.g.,
15 16	7	the death of a loved one or loss of a job), and the ERS enabled a regaining of structure and
17 18	8	control. Strugglers perceived the scheme as a way to get lives 'back on track' (Margaret).
19 20	9	This, when combined with complex health problems, meant disengagement could be difficult.
21 22 23	10	Brian, a widower with depression and a history of myocardial infarction, had: 'trouble with
23 24 25	11	my left foot, I am partially blind now, diabetic.' He reported a lack of confidence to move on,
26 27	12	indicating scheme dependency: 'I'm letting (staff member) set my programme. I might jigger
28 29	13	myself up. The scheme is fine. It's ideal. I've been asking him can I stop in it?' Margaret, in
30 31	14	contrast, felt her mental health had improved so she no longer needed to attend but
32 33	15	recognised that this was cyclical, and that she may need future support.
34 35	16	
36 37	17	Within struggle, participants reported increased PA and health gains 'I feel 100% better'
38 39 40	18	(Brian); however, longer-term positive continuation post-scheme appeared unlikely.
40 41 42	19	
43 44	20	Peter's experiences illustrate the theme of struggle. For him, social circumstance was
45 46	21	particularly influential. His attendance was sporadic (63% of potential sessions). He was
47 48	22	unemployed at his first interview and saw a potential return to work as an adherence barrier:
49 50	23	I juin a la companya de la companya
51 52		
53 54		
55 56 57		
57 58 59		
60		14 For peer review only - http://bmjopen.bmj.com/site/about/guidelines.xhtml

BMJ Open

1	Peter: Well I don't know exactly what is going to happen. I'll just see what it is and how it
2	goes. If I enjoy it, I will stick with it as long as I can. [] If I find work, I would stop
3	attending.'
4	
5	During his second interview, it was difficult to gauge his enthusiasm due to his natural
6	reticence: 'I come because it is there and it's available. Or I would be just moping around the
7	house all day doing nothing.' His intermittent attendance was due to a work-related course
8	and a short period of employment. Cost was an issue:
9	
10	Peter: 'It's pretty hard because I have got no wages coming in. I've just got Job Seekers
11	Allowance and that doesn't even pay my mortgage, so money is really tight. I did have a little
12	insurance but that money has now run out and I really am starting to struggle. It is fairly
13	cheap, but saying that, when you haven't got a lot of money coming in then it is a lot of
14	money to pay out.'
15	
16	Peter stated he would continue to attend until he found employment, when he 'might try and
17	come with my partner and play badminton once or twice a week.' His enthusiasm for this
18	appeared to be lacking and therefore long-term change in PA was unlikely.
19	
20	Defeat: inappropriate referral or poor participation experience
21	
22	The third theme was <i>defeat</i> . Within this, subthemes of poor health, social anxiety and poor
23	participation experience were apparent. Some participants never attended an exercise session,
24	being medically excluded (Paul), prevented by ill health (Dan) or social anxieties (despite
25	telephone support):
	15 For poor review only. http://breion.org/aits/about/avidalines.ubtral

1	
2	Jackie: 'The thought of coming here on my own, with nobody else, I like staying in my
3	comfort zone '(staff) phoned; she says about the sessions and that and I was being
4	honest with her so she left it a couple of weeks and then phoned back and she says would
5	you not like to come along by yourself? And I went no'
6	
7	For others, poor participation experience (Dorothy) indicated weaknesses in scheme delivery.
8	This group were defeated by the barriers faced, they felt ostracised from participation. There
9	was a sense of failure and, for some, shame. For these participants the ERS did not work.
10	
11	Illustrating this theme is Dorothy, who had been previously referred to the scheme on two
12	occasions. The first time she completed the scheme but did not continue via signposted exit
13	route activities because her friend stopped attending. On the second occasion, she dropped
14	out due to a foot problem. She did not like PA:
15	
15 16	Dorothy: 'At school I wasn't very good at PE. I never liked it very much, only did what I had
	Dorothy: 'At school I wasn't very good at PE. I never liked it very much, only did what I had to do. [] I've been to be the scheme twice before; other than that I don't think I really did
16	
16 17	to do. [] I've been to be the scheme twice before; other than that I don't think I really did
16 17 18	to do. [] I've been to be the scheme twice before; other than that I don't think I really did any exercise. [] I've got a real problem with my back. I'm sort of hoping that if I do exercise
16 17 18 19	to do. [] I've been to be the scheme twice before; other than that I don't think I really did any exercise. [] I've got a real problem with my back. I'm sort of hoping that if I do exercise
16 17 18 19 20	to do. [] I've been to be the scheme twice before; other than that I don't think I really did any exercise. [] I've got a real problem with my back. I'm sort of hoping that if I do exercise it will strengthen the muscles in my back and I will be able to do more things.'
16 17 18 19 20 21	to do. [] I've been to be the scheme twice before; other than that I don't think I really did any exercise. [] I've got a real problem with my back. I'm sort of hoping that if I do exercise it will strengthen the muscles in my back and I will be able to do more things.' During her second interview, Dorothy described how she felt unable to cope with the sessions
16 17 18 19 20 21 22	to do. [] I've been to be the scheme twice before; other than that I don't think I really did any exercise. [] I've got a real problem with my back. I'm sort of hoping that if I do exercise it will strengthen the muscles in my back and I will be able to do more things.' During her second interview, Dorothy described how she felt unable to cope with the sessions
16 17 18 19 20 21 22 23	to do. [] I've been to be the scheme twice before; other than that I don't think I really did any exercise. [] I've got a real problem with my back. I'm sort of hoping that if I do exercise it will strengthen the muscles in my back and I will be able to do more things.' During her second interview, Dorothy described how she felt unable to cope with the sessions due to back pain:
16 17 18 19 20 21 22 23 24	 to do. [] I've been to be the scheme twice before; other than that I don't think I really did any exercise. [] I've got a real problem with my back. I'm sort of hoping that if I do exercise it will strengthen the muscles in my back and I will be able to do more things.' During her second interview, Dorothy described how she felt unable to cope with the sessions due to back pain: Dorothy: 'I did tell her that I'd got a back problem and I was waiting for these injections, but

BMJ Open

1	was asking us to do. It was a class and it was where you do sort of aerobics first and then go
2	to all these sort of stations. I found it really hard.'
3	
4	Discussing one session, she described a lack of staff support: 'She just said do what you can
5	do and if you can't do whatever it is, just keep your feet moving.' This contrasted with her
6	experience with another staff member: 'She knew when you couldn't do it and she would give
7	you an alternative. So she was really good but she was only there once.'
8	
9	The scheme had a system of telephone support, but in Dorothy's case, implementation
10	appeared to be lacking:
11	
12	Dorothy: 'I 'phoned in several times to explain. I left messages but nobody got back to me. I
13	think if maybe someone had 'phoned me back and said 'well come in and you can do the
14	things a different way' it might have encouraged me to go back in again.'
14 15	things a different way' it might have encouraged me to go back in again.'
	things a different way' it might have encouraged me to go back in again.' Dorothy raised delivery issues and highlighted the need for a better understanding of protocol
15	
15 16	Dorothy raised delivery issues and highlighted the need for a better understanding of protocol
15 16 17	Dorothy raised delivery issues and highlighted the need for a better understanding of protocol implementation by staff. She was very upset by her experience, stating <i>'It makes you feel like</i>
15 16 17 18	Dorothy raised delivery issues and highlighted the need for a better understanding of protocol implementation by staff. She was very upset by her experience, stating ' <i>It makes you feel like a failure sometimes</i> . <i>I don't think I could go to the doctors and say 'I failed last time can you</i>
15 16 17 18 19	Dorothy raised delivery issues and highlighted the need for a better understanding of protocol implementation by staff. She was very upset by her experience, stating 'It makes you feel like a failure sometimes. I don't think I could go to the doctors and say 'I failed last time can you refer me again'?' Her experience was complex however. She felt unable to access the peer
15 16 17 18 19 20	Dorothy raised delivery issues and highlighted the need for a better understanding of protocol implementation by staff. She was very upset by her experience, stating ' <i>It makes you feel like a failure sometimes</i> . <i>I don't think I could go to the doctors and say 'I failed last time can you refer me again'?</i> ' Her experience was complex however. She felt unable to access the peer support described by those who engaged successfully. During her first interview, Dorothy
15 16 17 18 19 20 21	Dorothy raised delivery issues and highlighted the need for a better understanding of protocol implementation by staff. She was very upset by her experience, stating ' <i>It makes you feel like a failure sometimes</i> . <i>I don't think I could go to the doctors and say 'I failed last time can you refer me again'?</i> ' Her experience was complex however. She felt unable to access the peer support described by those who engaged successfully. During her first interview, Dorothy described how she found socialising difficult: ' <i>I'm not really a good mixer so I find it quite</i>
15 16 17 18 19 20 21 22	Dorothy raised delivery issues and highlighted the need for a better understanding of protocol implementation by staff. She was very upset by her experience, stating 'It makes you feel like a failure sometimes. I don't think I could go to the doctors and say 'I failed last time can you refer me again'?' Her experience was complex however. She felt unable to access the peer support described by those who engaged successfully. During her first interview, Dorothy described how she found socialising difficult: 'I'm not really a good mixer so I find it quite hard.' During her second interview, she reported that the group felt unwelcoming: 'She was
15 16 17 18 19 20 21 22 23	Dorothy raised delivery issues and highlighted the need for a better understanding of protocol implementation by staff. She was very upset by her experience, stating 'It makes you feel like a failure sometimes. I don't think I could go to the doctors and say 'I failed last time can you refer me again'?' Her experience was complex however. She felt unable to access the peer support described by those who engaged successfully. During her first interview, Dorothy described how she found socialising difficult: 'I'm not really a good mixer so I find it quite hard.' During her second interview, she reported that the group felt unwelcoming: 'She was really nice, one lady that was there. Apart from her, I don't think any of the others were

Lack of social confidence contributed to dropout, although in Dorothy's case it does not
 appear to be have been the primary influence. Defeat illustrates how some participants may
 struggle to access the peer support identified by others as an important facilitator for
 adherence.

DISCUSSION

The purpose of this study was to understand experiences of an ERS to give insight regarding what worked, or did not work, to encourage engagement, and for whom. This is important because existing literature questions ERS effectiveness,^{20 22} without exploring adequately how to focus implementation to better personalise support. Three overarching themes emerged. First, success, with engaged participants focused on health outcomes. Second, struggle, short-term success but with concerns regarding continued engagement. Finally, *defeat*, where illness, social anxiety, and/or poor participation experience prevented engagement. Within the identified themes, similarities in factors affecting engagement and non-engagement were evident. For those who experienced some measure of success, there were shared enablers such as peer support, achievable and enjoyable activities, staff knowledge and staff/peer support. These are reflected to some extent in systematic review findings of engagement facilitators for ERS.³¹ What this paper adds is insight regarding whom these facilitators worked best for within the ERS context. Specifically, participants who were able to access social support, had positive previous experiences of PA, and were motivated by improving or maintaining their health. Participants often described success in terms of improvements in mental health and self-esteem, perceptions about ability to be active, and the social benefits of participation. Social and psychological benefits were perceived to be as meaningful as measurable physical health benefits, similar to other

BMJ Open

reported findings.³² Success illustrated the value of exercise referral for some participants
with non-communicable diseases. This was not universal however. The study highlighted
unequal abilities to access the scheme, along with differing support requirements, which
suggests the need to provide more tailored support for some. The issues identified are
reflective of other studies examining barriers to PA irrespective of the presence of a medical
condition.⁷¹¹

Adults with complex lives embarked on the ERS with expectations of positive changes in health. While ERS delivery training courses include elements of behaviour change, the training does not appear to be sufficient preparation for staff to deal with identified complex psychological barriers. Indeed the high levels of responsibilities that fitness professionals undertake has led to concerns about adequacy of education and training.³³ Further development of behaviour change elements within national occupational standards for promoting PA could partially address matters. This represents only part of the problem however. This study illustrates how a 'one size fits all' model does not adequately cater for the complex range of referrals received. Indeed, the existing model of universal referral to a common programme is potentially setting such schemes up to fail. This is because current measures of success are typically quantified as uptake and adherence,³⁴⁻³⁶ and/or self-reported changes in PA.^{15 37} Regardless of suitability, providers may feel obliged to 'shoehorn' referrals into schemes if the continuation of funding is reliant on achievement of such key performance indicators. This type of approach fails to consider the complex health and social circumstances of ERS participants, leading to an inadequate focus on what works, and for whom.

1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
11	
12	
13	
14	
15	
16	
17	
18	
19 20	
20	
21	
22	
23	
24	
25	
26	
27	
28	
29	
30	
31	
32	
33	
34	
35	
36	
37	
38	
39	
40	
40 41	
42 43	
44 45	
46	
47	
48	
49	
50	
51	
52	
53	
54	
55	
56	
57	
58	
59	
60	

1

1	In the case of ERS, there is a need to understand what different approaches are required to
2	support change for those experiencing struggle or defeat. The themes presented in this study
3	may resonate with ERS commissioners and providers and should encourage reflection of
4	approaches to support. Success can reinforce good practice, while highlighting potential
5	improvements. Struggle can initiate conversations about alternative delivery for those who
6	require more or different support in order to make sustained behaviour change. This may
7	include mechanisms for cyclical support to reengage those who relapse into inactivity and
8	'weaning' to reduce ERS dependency. Finally, defeat can initiate conversations about
9	appropriate referrals, improvements to existing provision and alternative models of care. At a
10	broad level, approaches may include support from multiple agencies, ³⁸ the use of
11	technology, ³⁹ or broader system change. ⁴⁰ Promisingly, there is emerging evidence of
12	practice with the potential to better support patients with struggle or defeat-style narratives.
13	Those with poor health may benefit from individualisation of exercise, ⁴¹ those with social
14	anxiety from more online delivery and support, ⁴² and the complex needs of patients are more
15	likely to be catered for appropriately with increasing use of scheme co-production. ⁴³ Calls at
16	national policy level for better use of triage or a 'stepped approach to delivery'21 may further
17	assist with both enhancing support for those with challenging circumstances and modifying
18	or reducing it for those that risk becoming scheme dependent. Testing the effectiveness of
19	these ideas should be a priority for future research.
20	

- 21 Methodological considerations
- 22
- 23

Qualitative analysis is inherently subjective since it is influenced by the assumptions, beliefs
and biases of the researcher.⁴⁴ In this case, the researcher was experienced in the management

BMJ Open

3	
4	
5	
6	
7	
8	
9	
10	
11	
12	
13	
14	
15	
16	
17	
18	
19	
20	
21	
22	
23	
24	
25	
26	
27	
28	
29	
30	
31	
32	
33	
34	
35	
36	
37	
38 39	
39 40	
40 41	
41 42	
42 43	
44	
45	
46	
47	
48	
49	
50	
51	
52	
53	
54	
55	
56	
57	
58	
59	
60	

1 and delivery of the ERS studied. Potential biases were explored by the use of reflective field 2 notes and in group discussions with all authors. Particular attention was paid to how existing 3 knowledge may have affected discussion with participants and interpretation of results. That 4 said, while in the past an outsider, objective stance was considered desirable in research terms to guard against identification, insider insight can now be considered legitimate and desirable 5 due to the potential for increased empathy with participants.⁴⁵ After reflection, it was felt that 6 researcher knowledge contributed positively to the interpretation of data through being able 7 to understand the particular scheme that participants were discussing. 8 9 10 For each participant, interviews took place on two occasions. Qualitative interviews are only 11 able to uncover what participants recall or are willing to reveal about their experiences at a 12 particular time, rather than realities. As such they may reflect recall bias or inaccuracies. 13 Participant knowledge of the researcher background may also have influenced what was disclosed. Readers can make choices about whether the identified themes resonate with their 14 15 own intuitive understanding of such situations, which arguably can improve practice through the process of naturalistic generalization.⁴⁶ 16 17 18 It is not known whether the experiences of those who declined to participate or dropped out 19 of the study were different to those who took part. For example, we previously established that those under 55 years of age were less likely to engage in the first instance and more 20 likely to dropout when they did.²⁴ However, only one participant from this demographic 21 22 completed a second interview. Additionally, this piece of work did not examine barriers to 23 scheme access for who did not attend the initial consultation. Understanding this group, 24 however, is critical for determining who current services are failing and why. 25

1 CONCLUSION

3	Overall, the data support arguments that ERSs can disproportionally engage with, and
4	benefit, some disadvantaged groups. Importantly they can successfully engage those with
5	non-communicable diseases, and positively affect health and wellbeing. The value of current
6	ERSs appears to be for those with social confidence and previous positive experiences of PA.
7	Conversely, such schemes may fail those who struggle to access social support due to varying
8	health condition demands, or complex or impaired social circumstances. For those who are
9	unable to adhere, feelings of ostracism and failure may further exacerbate outcome
10	differentials. Ultimately, even programmes that target disadvantaged sub-groups (in the case
11	of ERSs, those with non-communicable diseases) appear at risk of reinforcing inequalities.
12	This study therefore highlights a need for services and systems that better provide for those
13	with dynamic health and social circumstances.
14	
15	Acknowledgements: Many thanks to ERS staff at Blyth Valley Arts and Leisure for helping
16	with recruitment and organisation of interviews.
17	
18	Author Contributions: CLH contributed to the study design, data collection, data analysis
19	and preparation of the final document. EO contributed to data analysis and preparation of the
20	final document. CDR contributed to study design, data analysis and preparation of the final
21	document. LJA contributed to study design, data analysis and preparation of the final
22	document. All authors contributed to this paper and approved the final version.
23	
24	Funding: Blyth Valley Arts and Leisure was one of two providers of the exercise referral
25	scheme studied and provided funding for a PhD to evaluate the service.

BMJ Open

2 3	1	
4		
5 6	2	Competing interests: CLH is a former employee of Blyth Valley Arts and Leisure and
7 8	3	completed a PhD that was funded by the aforementioned company.
9 10	4	
11	5	Participant consent: obtained
12 13	5	
14 15	6	
15 16 17	7	Ethics approval: Northumbria University Faculty of Health and Life Sciences Ethics
18 19	8	Committee (Ref: 15-03-131781)
20 21	9	
22	10	Provenance and peer review: Not commissioned; externally peer reviewed.
23 24	11	
25 26		
27 28	12	Data sharing statement: No additional data are available.
29	13	
30 31 32	14	REFERENCES
33 34	15	REFERENCES
35 36	16	1. Lee IM, Shiroma EJ, Lobelo F, et al. Effect of physical inactivity on major non-
37 38	17	communicable diseases worldwide: an analysis of burden of disease and life expectancy.
39 40 41	18	Lancet 2012;380:219-29. doi: 10.1016/S0140-6736(12)61031-9
42 43	19	2. Ding D, Lawson KD, Kolbe-Alexander TL, et al. The economic burden of physical
44 45	20	inactivity: a global analysis of major non-communicable diseases. Lancet 2016;388:1311-24.
46 47 48	21	doi: 10.1016/S0140-6736(16)30383-X
49 50	22	3. World Health Organisation. Global action plan for the prevention and control of non-
51 52	23	communicable diseases 2013-2020. Geneva: World Health Organisation, 2013.
53 54	24	http://africahealthforum.afro.who.int/IMG/pdf/global_action_plan_for_the_prevention_and_c
55 56 57	25	ontrol_of_ncds_2013-2020.pdf: (Accessed: 05 March 2018)
58 59 60		23 For peer review only - http://bmjopen.bmj.com/site/about/guidelines.xhtml

1	4. World Health Organisation. Global status report on noncommunicable diseases, 2014.
2	Geneva: World Health Organisation, 2014.
3	http://apps.who.int/iris/bitstream/10665/148114/1/9789241564854_eng.pdf (Accessed: 05
4	March 2018).
5	5. Eime RM, Charity MJ, Harvey JT, et al. Participation in sport and physical activity:
6	associations with socio-economic status and geographical remoteness. BMC Public Health
7	2015;15:434. doi: 10.1186/s12889-015-1796-0
8	6. Sievi NA, Senn O, Brack T, et al. Impact of comorbidities on physical activity in COPD.
9	Respirology 2015;20(3):413-18. doi: 10.1111/resp.12456
10	7. Picorelli AMA, Pereira LSM, Pereira DS, et al. Adherence to exercise programs for older
11	people is influenced by program characteristics and personal factors: a systematic review.
12	Journal of Physiotherapy 2014;60(3):151-56. doi: 10.1037/0022-006X.51.3.390
13	8. Allender S, Cowburn G, Foster C. Understanding participation in sport and physical
14	activity among children and adults: a review of qualitative studies. Health Educ Res
15	2006;21(6):826-35. doi: 10.1093/her/cyl063
16	9. Borodulin K, Sipila N, Rahkonen O, et al. Socio-demographic and behavioral variation in
17	barriers to leisure-time physical activity. Scand J Public Health 2016;44(1):62-69. doi:
18	10.1177/1403494815604080
19	10. Kern ML, Reynolds CA, Friedman HS. Predictors of Physical Activity Patterns Across
20	Adulthood: A Growth Curve Analysis. Pers Soc Psychol Bulletin 2010. doi:
21	10.1177/0146167210374834

BMJ Open

1	11. Pan S, Cameron C, DesMeules M, et al. Individual, social, environmental, and physical
2	environmental correlates with physical activity among Canadians: a cross-sectional study.
3	BMC Public Health 2009;9(1):21. doi: 10.1186/1471-2458-9-21
4	12. Zunft H-JF, Friebe D, Seppelt B, et al. Perceived benefits and barriers to physical activity
5	in a nationally representative sample in the European Union. Public Health Nutr
6	1999;2(Supplement 1a):153-60. doi: 10.1017/S1368980099000208
7	13. Sowden S, Raine R. Running along parallel lines: how political reality impedes the
8	evaluation of public health interventions. A case study of exercise referral schemes in
9	England. J Epidemiol Community Health 2008;62(9):835 - 41. doi:10.1136/jech.2007.069781
10	14. Murphy SM, Edwards RT, Williams N, et al. An evaluation of the effectiveness and cost
11	effectiveness of the National Exercise Referral Scheme in Wales, UK: a randomised
12	controlled trial of a public health policy initiative. J Epidemiol Community Health
13	2012;66(8):745-53. doi: 10.1136/jech-2011-200689
14	15. Sorenson J, Kragstrup J, Stovgaard T, et al. Exercise on Prescription: a randomized study
15	on the effect of counseling vs counseling and supervised exercise. Scand J Med Sci Sports
16	2008;18(3). doi: 10.1111/j.1600-0838.2008.00811.x
17	16. Gusi N, Reyes M, Gonzalez-Guerrero J, et al. Cost-utility of a walking programme for
18	moderately depressed, obese, or overweight elderly women in primary care: a randomised
19	controlled trial. BMC Public Health 2008;8(1):231. doi: 10.1186/1471-2458-8-231
20	17. Gallegos-Carrillo K, García-Peña C, Salmerón J, et al. Brief Counseling and Exercise
21	Referral Scheme: A Pragmatic Trial in Mexico. Am J Prev Med 2017;52(2):249-59.
22	doi:10.1016/j.amepre.2016.10.021

	1	18. Public Health England. Identifying what works for local physical inactivity interventions.
	2	London, Public Health England, 2014.
	3	https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/374560/What
	4	worksv1_2.pdf: (Accessed: 05 March 2018)
	5	19. Oliver EJ, Hanson CL, Lindsey IA, et al. Exercise on referral: evidence and complexity at
	6	the nexus of public health and sport policy. International Journal of Sport Policy and Politics
	7	2016:1-6. doi: 10.1186/1471-2458-9-21
	8	20. Campbell F, Holmes M, Everson-Hock E, et al. A systematic review and economic
	9	evaluation of exercise referral schemes in primary care: a short report. Health Technol Assess
1	10	2015;19(60):1-87. doi: 10.3310/hta19600
1	11	21. National Institute for Health and Care Excellence. Physical activity; exercise referral
1	12	schemes. National Institute for Health and Care Excellence, 2014.
1	13	https://www.nice.org.uk/guidance/ph54/resources/physical-activity-exercise-referral-
1	14	schemes-pdf-1996418406085 (Accessed: 05 March2018).
1	15	22. Pavey TG, Taylor A, Fox K, et al. Effect of exercise referral schemes in primary care on
1	16	physical activity and improving health outcomes: systematic review and meta-analysis. BMJ
1	17	2011;343. doi: 10.1136/bmj.d6462
1	18	23. Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research
1	19	(COREQ): a 32-item checklist for interviews and focus groups. Int J Qual Health Care
2	20	2007;19(6):349-57. doi: 10.1093/intqhc/mzm042
2	21	24. Hanson CL, Allin LJ, Ellis JG, et al. An evaluation of the efficacy of the exercise on
2	22	referral scheme in Northumberland, UK: association with physical activity and predictors of
2	23	engagement. A naturalistic observation study. BMJ Open 2013;3(8). doi: 10.1136/bmjopen-
2	24	2013-002849
		26 For peer review only - http://bmjopen.bmj.com/site/about/guidelines.xhtml

BMJ Open

1	25. Prochaska JO, Marcus BH. The transtheoretical model: Applications to exercise. In:
2	Dishman RK, ed. Advances in exercise adherence. Champaign, IL, England: Human Kinetics
3	1994:161-80.
4	26. Prochaska JO, DiClemente CC. Stages and processes of self-change of smoking: Toward
5	an integrative model of change. J Consult and Clin Psychol 1983; Vol 51(3):390-95. doi:
6	10.1037/0022-006X.51.3.390
7	27. Marshall MN. Sampling for qualitative research. Fam Pract 1996;13(6):522-26.
8	doi:10.1093/fampra/13.6.522
9	28. Strauss AL. Qualitative analysis for social scientists. Cambridge: Cambridge University
10	Press, 1987.
11	29. Spencer L, Ritchie J, Lewis J, et al. Quality in qualitative evaluation: a framework for
12	assessing research evidence. National Centre for Social Research, 2003.
13	http://dera.ioe.ac.uk/21069/2/a-quality-framework-tcm6-38740.pdf (Accessed: 05 March
14	2018).
15	30. Festinger L. A Theory of Social Comparison Processes. Human Relations 1954;7(2):117-
16	40. doi: 10.1177/001872675400700202
17	31. Morgan F, Battersby A, Weightman AL, et al. Adherence to exercise referral schemes by
18	participants - what do providers and commissioners need to know? A systematic review of
19	barriers and facilitators. BMC Public Health 2016;16(1):1-11 doi: 10.1186/s12889-016-2882-
20	7
21	32. Pentecost C, Taket A. Understanding exercise uptake and adherence for people with
22	chronic conditions: a new model demonstrating the importance of exercise identity, benefits
23	of attending and support. Health Educ Res 2011;26(5):908-22. doi: 10.1093/her/cyr052
	27 For recer review only http://hypeionen.htmi.com/cite/ahout/avidalines.yhtml

For peer review only - http://bmjopen.bmj.com/site/about/guidelines.xhtml

1	33. De Lyon ATC, Neville RD, Armour KM. The Role of Fitness Professionals in Public
2	Health: A Review of the Literature. Quest 2017;69(3):313-30. doi:
3	10.1080/00336297.2016.1224193
4	34. Dugdill L, Graham R, McNair F. Exercise referral: the public health panacea for physical
5	activity promotion? A critical perspective of exercise referral schemes; their development and
6	evaluation. Ergonomics 2005;48:1390-410. doi: 10.1080/00140130500101544
7	35. Gidlow C, Johnston LH, Crone D, et al. Socio-demographic patterning of referral, uptake
8	and attendance in Physical Activity Referral Schemes. J Public Health 2007;29(2):107-13.
9	doi:10.1093/pubmed/fdm002
10	36. Sowden S, Breeze E, Barber J, et al. Do general practices provide equitable access to
11	physical activity interventions? Br J Gen Pract 2008;58:e1-e8. doi: 10.3399/bjgp08X342237
12	37. Duda JL, Williams GC, Ntoumanis N, et al. Effects of a standard provision versus an
13	autonomy supportive exercise referral programme on physical activity, quality of life and
14	well-being indicators: a cluster randomised controlled trial. Int J Behav Nutr Phys Act
15	2014;11(1):1-15. doi: 10.1186/1479-5868-11-10
16	38. Sugai G, Horner RH, Dunlap G, et al. Applying positive behavior support and functional
17	behavioral assessment in schools. J Posit Behav Interv 2000;2(3):131-43. doi:
18	10.1177/109830070000200302
19	39. Gallagher R, Gallagher P, Zhang L, et al. Technology Use and Confidence in Cardiac
20	Rehabilitation Patients. Heart Lung Circ 2016;25:S333. doi: 10.1016/j.hlc.2016.06.791
21	40. Michie S, van Stralen MM, West R. The behaviour change wheel: A new method for
22	characterising and designing behaviour change interventions. Implement Sci 2011;6(1):42.
23	doi: 10.1186/1748-5908-6-42

Page 29 of 41

BMJ Open

1	41. Martín-Borràs C, Giné-Garriga M, Puig-Ribera A, Martín C, Solà M, Cuesta-Vargas AI.
2	A new model of exercise referral scheme in primary care: is the effect on adherence to
3	physical activity sustainable in the long term? A 15-month randomised controlled trial. BMJ
4	open. 2018;8(3):e017211.
5	42. Hawkins J, Madden K, Fletcher A, Midgley L, Grant A, Cox G, et al. Development of a
6	framework for the co-production and prototyping of public health interventions. BMC Public
7	Health. 2017;17:689.
8	43. Buckley B, Thijssen D, Murphy R, Graves L, Whyte G, Gillison F, et al. Making a move
9	in exercise referral: co-development of a physical activity referral scheme. Journal of Public
10	Health. 2018. doi: 0.1093/pubmed/fdy072
11	44. Creswell JW, Miller DL. Determining validity in qualitative inquiry. <i>Theory into practice</i>
12	2000;39(3):124-30. doi: 10.1207/s15430421tip3903
13	45. Gair S. Feeling Their Stories: Contemplating Empathy, Insider/Outsider Positionings, and
14	Enriching Qualitative Research. Qual Health Res 2012;22(1):134-43. doi:
15	10.1177/1049732311420580
16	46. Stake RE, Trumbull DJ. Naturalistic generalizations. Review Journal of Philosophy and
17	Social Science 1982;7(1):1-12.
18	

BMJ Open Referral by healthcare professional (Standardised referral form)

Personal contact from ERS

(Telephone call or postal contact to arrange initial one to one consultation)

Initial consultation (All consultations based on Transtheoretical Model)

Assessment of stage of change, assessment of self-reported physical activity and discussion about physical activity levels / opportunities to exercise via the scheme (behaviour change techniques such as decisional balance incorporated as required) Not ready to change or medically excluded

Brief physical activity advice notification to referrer of nonparticipation

Ready to change

Supervised ERS group physical activity sessions 12 weeks, 2 sessions per week Options include gym usage, circuit classes, racquet sports and aqua aerobics

Individual non-ERS supervised physical activity options Swimming, casual gym or fitness class attendance

12-week consultation

Assessment of stage of change, discussion about physical activity levels / opportunities to exercise via the scheme (behaviour change techniques incorporated as required) Non-attenders return encouraged via telephone (3 attempts) / postal contact (if unable to contact by telephone

Supervised ERS group physical activity sessions 12 weeks, 2 sessions per week Options include gym usage, circuit classes, racquet sports and aqua aerobics

Individual non-ERS supervised physical activity options Swimming, casual gym or fitness class attendance

24-week consultation

Assessment of stage of change, assessment of self-reported physical activity and discussion about physical activity levels / opportunities to exercise after completion of the scheme (behaviour change techniques incorporated as required) Non-attenders return encouraged via telephone (3 attempts) / postal contact (if unable to contact by telephone

Feedback to referrer

Exit routes (similar ERS supervised exit sessions, reduced cost fitness and/or swimming memberships) or independent activity

For peer review only - http://bmjopen.bmj.com/site/about/guidelines.xhtml




Figure 2: Thematic analysis of ERS experiences

206x148mm (300 x 300 DPI)

Expectations of participants in the Northumberland exercise on referral scheme.

Preface: Set the interviewee at rest; explain the purpose of the interview (to understand their expectations of the exercise on referral scheme) explain that the expected outcomes (that the study will give a better understanding of why the scheme works for some people, but not others); rules of confidentiality etc.

Record demographic / personal info to start:

- Gender
- Age

 • Employment status

Question	Prompts
1. Tell me a bit about what sort of physical activity you have taken part in in the past.	 What were your experiences of sport /physical activity at school / as a child What influence have others had on the type of physical activity you have taken part in? Do you have any particular likes / dislikes of sport / physical activity? Tell me about any times in past where there has been a big change in your physical activity patterns Has there been anything else that has influenced your participation in physical activity?
2. How do you feel about taking part in physical activity now?	 What type of physical activity (if any) do they take part in at the moment? What type of physical activity would they like to take part in? What do they think the important reasons for taking part in physical activity are? Is there anything that particularly worries them about taking part in physical activity?
3. So thinking about the exercise on referral scheme that you have been referred to, how did you find out about it?	 Who/what has motivated you to attend? What made you decide that this is the right time to take part in the scheme?
4. Why were you referred to the scheme?	 What did the referrer explain to you about the scheme? What do you expect (if anything) when you start attending the scheme?

 BMJ Open

 What are the changes to your health that you expect will happen as a result of participation? How quickly do you expect to see these changes? How have you decided that these changes are realistic? How confident do you feel about taking part in the scheme?
 How quickly do you expect to see these changes? How have you decided that these changes are realistic?
• How have you decided that these changes are realistic?
How confident do you feel about taking part in the scheme?
What are you particularly looking forward to?
• What are you worried about?
• How long after referral did it take to be contacted by the scheme?
• What information has been given to you prior to the initial consultation?
• How comfortable do you feel coming to first consultation?
CL:
How important do you think attendance in a group will be?
• What do you expect from the staff on the scheme?
How important are changes in health?
Why were the influences raised important?
• Tell me about any worries you might have about health issues
• Tell me about any other things such as other commitments that might stop you from
attending
• What ways might you overcome these issues if they arise?
like to tell me about your expectations for participation in the scheme?
-

Semi Structured Interview Questions:

 Welcome interviewee back; explain the purpose of the interview (to hear about their experiences of the exercise on referral scheme) explain that the expected outcomes (that the study will give a better understanding of why the scheme works for some people, but not others); rules of confidentiality etc.

1. First I would like to	• What did the member of staff ask you about?(medical issues related to exercise, past and current exercise, feelings
talk about the	about taking part in the scheme)
consultation that	• How did these questions make you feel?
you had before you	• What tests were carried out?
started the scheme -	
what you were	test)?
asked about and	 What information were you given? (scheme information, times of sessions etc, results of tests that were carried
what information	out, other information about physical activity options outside the scheme, cost of attendance)
you were given.	 Did you agree to attend at the end of the consultation?(if no go on to question 2)
	 How did you feel about attending the scheme exercise sessions at the end of the consultation?
	 Was there anything that you were looking forward to?
	• Was there anything that you were concerned about
2. Why did you decide	How did you feel about being referred at the end of the consultation?
that the scheme was	• Did the consultation encourage you to take part in physical activity even though you did not attend any sessions?
not right for you?	If so, in what way?
	• Have you been back to see the person who referred you to discuss your referral? If so, what did you discuss?
	• What could the scheme have offered you that would have encouraged you to attend?
	• Have you changed your levels of physical activity since you were referred? If yes, did the fact that you were
	referred influence this and in what way
How many scheme exer	cise sessions have you attended?

 BMJ Open

3.	Thinking about how	• How comfortable did you feel about coming into the centre to take part in an actual exercise session?
	first exercise	What happened during the first session?
	session, how did you	• What activities did you take part in?
	feel about the	• How did you feel about the activities?
	scheme before you	• How did you find the other members of the group?
	attended this?	• How did you feel after the first session?
4.	Are you still	• What are the things that you enjoy about attending the scheme?
	attending the	• How have you found the staff?
	scheme sessions? (if	Have there been any weeks were you have not attended at all?
	no, go to question 6)	• If yes, did anyone from the scheme contact you?
		• If yes, who, how and did it have any effect?
		• Do you have any suggestions for what the scheme might do differently?(go to Q7)
5.	When did you stop	Why did you stop attending?
	attending?	• Did anyone from the scheme contact you when you did not attend?
		What happened?
		How did you find the staff?
		How did you find the sessions that you did attend?
		What, if any, parts of the scheme did you enjoy?
		• Do you have any suggestions for what the scheme might do differently?
		• Compared to before you were referred to the scheme, have you changed the amount of physical activity that you do?
		• If yes, what do you do that is different?
		What made you change your activity?

		• How important do you think that physical activity is in helping to maintain or improve your health? Has this view changed since you were referred?
6.	At the first interview, you told me that you hoped to achieve Now you have taken part in the scheme for 12 weeks:	 What changes to your health have you noticed? Were these changes what you expected? Have other factors outside the scheme had an effect on your health / lifestyle choices? How important are these changes in encouraging you to keep attending?
7.	At the first interview, you told me that you are worried aboutNow that you have taken part in the scheme for 12 weeks:	 Were the concerns that you had justified? How were these concerns addressed when you attended? Is there anything that you can suggest that the scheme might do to help other people who feel the same way abou attending as you did?
8.	Have you increased the amount of activity you do overall?	 Have you increased the amount of activity you are doing independently of the scheme sessions? If so, what you are doing that is different from before you started? Why have you changed your activity outside the scheme? How important do you think that physical activity is in maintaining or improving your health? Is this different to before you started?
9.	What are you most looking forward to in the next 12 weeks of the scheme?	 Is there anything in particular that you think will encourage you to keep attending? What are you hoping to achieve over the second part of the scheme?

BMJ Open

10. What things do you	• Tell me about any worries you might have about health issues and scheme attendance
	• Tell me about any other things such as other commitments that might stop you from attending
likely to prevent you	• Have you thought about any ways that you might overcome these issues if they arise?
from attending	
sessions?	
11. Is there anything else	that you would like to tell me about your experience of participation in the scheme and your expectations for the
rest of the scheme?	
	For peer review only
	For peer review only - http://bmjopen.bmj.com/site/about/guidelines.xhtml

COREQ GUIDELINES REPORTING CHECKLIST: How do participant experiences and characteristics influence engagement in exercise referral? A qualitative longitudinal study of a scheme in Northumberland, United Kingdom.

No	Item	Guide	Information	Reported in
		questions/description		manuscript (Section, page no)
	Domain 1. Rese	arch team and reflexivity		(Section, page no)
	Personal Charact	· · · · · · · · · · · · · · · · · · ·		
1	Interviewer/	Which author/s conducted	CLH	Data collection and
-	facilitator:	the interview or focus group?		analysis, page 9
2	Credentials	What were the researcher's credentials?	CLH PhD student and experienced in working in intervention studied. LJA experienced qualitative researcher	Data collection an analysis, page 9
3	Occupation	What was their occupation at the time of the study?	PhD student and strategic manager in the ERS provider organisation	Data collection an analysis, page 9
4	Gender	Was the researcher male or female?	Female (referred to as her)	Data collection an analysis, page 9
5	Experience and training	What experience or training did the researcher have?	CLH: Qualitative research training and support from experienced researcher (LJA)	Data collection an analysis, page 9
	Relationship with	n participants		
6	Relationship established	Was a relationship established prior to study commencement?	No, participants were approached by ERS staff to explain the study and seek consent.	Sample, page 8
7	Participant knowledge of the interviewer	What did the participants know about the researcher? <i>e.g. personal</i> <i>goals, reasons for doing</i> <i>the research</i>	Participants were informed of researcher's employment status and that the research aimed to improved service delivery	Sample, page 8
8	Interviewer characteristics	What characteristics were reported about the interviewer/facilitator? <i>e.g. Bias, assumptions,</i> <i>reasons and interests in</i> <i>the research topic</i>	Acknowledgement of potential bias due to insider knowledge. Interest in service improvement and employment status of interviewer disclosed in study invitation information.	Methodological Considerations, page 21-22 Sample, page 8
	Domain 1: Stud			
	Theoretical frame			
9	Methodological orientation and Theory	What methodological orientation was stated to underpin the study? <i>e.g.</i> <i>grounded theory</i> ,	Thematic analysis using the framework approach	Data collection an analysis, page 9

		discourse analysis, ethnography, phenomenology, content analysis		
	Participant select		1	1
10	Sampling	How were participants selected? <i>e.g. purposive,</i> <i>convenience, consecutive,</i> <i>snowball</i>	Initially convenience from a defined group, then purposive	Sample, page 8
11	Method of approach	How were participants approached? <i>e.g. face-to-</i> <i>face, telephone, mail,</i> <i>email</i>	By telephone by scheme provider	Sample, page 8
12	Sample size	How many participants were in the study?	15 initially but only 11 completed both interviews	Results, page 10
13	Non- participation	How many people refused to participate or dropped out? Reasons?	Of initial invitees 13/25 refused. Of 15 initial participants, 4 dropped out. (no response n=2, too busy n=2)	Sample, page 8 Results, page 10
	Setting			[
14	Setting of data collection	Where was the data collected? <i>e.g. home,</i> <i>clinic, workplace</i>	Leisure centre where ERS was attended	Data collection and analysis, page 9
15	Presence of non- participants	Was anyone else present besides the participants and researchers?	No, interviews were conducted in private	Data collection and analysis, page 9
16	Description of sample	What are the important characteristics of the sample? <i>e.g. demographic</i> <i>data, date</i>	Gender, age group, medical reason for referral, employment status and previous PA Date range for interviews reported	Results, Table 1 Data collection and analysis, page 9
	Data collection			
17	Interview guide	Were questions, prompts, guides provided by the authors? Was it pilot tested?	Semi-structured interview guide used. Pilot tested. Guides provided as supplementary files	Data collection and analysis, page 9 Supplementary file 1 and 2
18	Repeat interviews	Were repeat interviews carried out? If yes, how many?	Yes, one further interview after 12-20 weeks	Data collection and analysis, page 9
19	Audio/visual recording	Did the research use audio or visual recording to collect the data?	Yes, the interviews were audio recorded	Data collection and analysis, page 9
20	Field notes	Were field notes made during and/or after the interview or focus group?	Yes detailed field notes were made directly after interviews	Data collection and analysis, page 9
21	Duration	What was the duration of the interviews or focus groups?	Range and median length reported: 22-62 minutes (median 48 minutes)	Data collection and analysis, page 9
22	Data saturation	Was data saturation	Recruitment stopped	Sample, page 8

		discussed?	when no new themes overarching were emerging		
23	Transcripts returned	Were transcripts returned to participants for comment and/or correction?	Transcripts were checked by participants.	Data collection and analysis, page 9	
	Domain 3: analysis and findings Data analysis				
24	Number of data coders	How many data coders coded the data?	N=2 (CLH and LJA) independently N=4 in total at data workshops	Data collection an analysis, page 9	
25	Description of the coding tree	Did authors provide a description of the coding tree?	Yes figure 2 visually describes the coding tree	Figure 2	
26	Derivation of themes	Were themes identified in advance or derived from the data?	Identified from data	Data collection an analysis, page 9-1	
27	Software	What software, if applicable, was used to manage the data?	No software used, data analysed manually using an excel spreadsheet	Data collection an analysis, page 9	
28	Participant checking	Did participants provide feedback on the findings?	No, but themes were checked with ERS staff at a workshop.	Data collection an analysis, page 10	
	Reporting				
29	Quotations presented	Were participant quotations presented to illustrate the themes / findings? Was each quotation identified? e.g. participant number	Yes, participants identified using a pseudonym	Results, page 10-1	
30	Data and findings consistent	Was there consistency between the data presented and the findings?	Themes were illustrated by participant quotations	Results, page 10-1	
31	Clarity of major themes	Were major themes clearly presented in the findings?	Three major experience themes were identified: success, struggle and defeat	Results, page 10-1	
32	Clarity of minor themes	Is there a description of diverse cases or discussion of minor themes?	Minor themes identified within each major theme: For success: improved health, increased PA, enjoyment and support For struggle: scheme dependency, multiple barriers and cyclical needs For defeat: inappropriate referral, poor participation experience, social anxiety.	Results, page 10-1	

Page 41 of 41

2	
2 3	Diverse cases discussed
4	within themes
5 6	
7	
8 9	
9 10	
11	
12	
13	
14	
15	
16 17	
17 18	
19	
20	
21	
22	
23 24	
24 25	
26	
27	
28	
29	
30 31	
32	
33	
34	
35	
36 37	
38	
39	
40	
41	
42	
43 44	
45	
46	
47	
48	
49 50	
50 51	
52	
53	
54	
55	
56 57	
57 58	
59	
60	For peer review only - http://bmjopen.bmj.com/site/about/guidelines.xhtml