

Oxyphor 2P: A High-Performance Probe for Deep-Tissue Longitudinal Oxygen Imaging

METHODS S1

NMR and MALDI-TOF spectra

Related to STAR Methods

Tatiana V. Esipova,^{1,5} Matthew J. P. Barrett,^{3,5} Eva Erlebach,³ Artëm E. Masunov,⁴ Bruno Weber*³ and
Sergei A. Vinogradov*^{1,2,6}

¹ Department of Biochemistry and Biophysics, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104, USA.

² Department of Chemistry, School of Arts and Sciences, University of Pennsylvania, Philadelphia, PA 19104, USA.

³ Institute of Pharmacology and Toxicology, University of Zurich, and Neuroscience Center Zurich, Zurich CH-8057, Switzerland.


⁴ NanoScience Technology Center, Department of Chemistry and School of Modeling, Simulation and Training, University of Central Florida, Orlando, FL 32826, USA; National Research Nuclear University MEPhI, Kashirskoye Shosse 31, Moscow 115409, Russia; South Ural State University, Lenin Pr. 76, Chelyabinsk 454080, Russia.

⁵ These authors contributed equally to this work.


⁶ Lead contact.

Correspondence: vinograd.upenn@gmail.com, bweber@pharma.uzh.ch



1: ¹H NMR





2: ^{13}C NMR




3: ^1H NMR





4: ^1H NMR






4: ^{13}C NMR




5: MALDI-TOF



5: ^1H NMR



6: MALDI-TOF



Oxyphor 2P: MALDI-TOF

