

Construction of a high-density linkage map and QTL mapping for important agronomic traits in *Stylosanthes guianensis* (Aubl.)

Sw.

Yan-Qiong Tang^{1*}, Zhi-Qiang Xia^{2*} Ze-Ting Ding^{1*}, Ya-Cao Ding¹, Zhu Liu¹, Xiang Ma¹ &
Jin-Ping Liu¹

¹Hainan Key Laboratory for Sustainable Utilization of Tropical Bioresources, Tropical Agriculture and Forestry Institute, Hainan University, Haikou, Hainan Province 570228, China.²The Institute of Tropical Bioscience and Biotechnology, Chinese Academy of Tropical Agricultural Sciences, Haikou, Hainan Province 571101, China

Correspondence and requests for materials should be addressed to Y.-Q. T. (tyq68@126.com) or J.-P. L. (liu3305602@163.com)

* These authors contributed equally to this work.

Table S3. Phenotype variations of the yield-related traits for F₂ population derived from the cross between TPRC1979 (female) and TPRCR273 (male) of *Stylosanthes guianensis*. CV: Coefficient of variation; FW: fresh weight; DW: dry weight; FW/DW: the fresh weight: dry weight ratio; PH: natural height; PBN: primary branch number; PW: plant width; LB: the maximum length of branches; LL: leaf length; LW: leaf width; LL/LW: the leaf length: leaf width ratio.

Traits	Mean	St. D	Min	Max	Range	CV(%)
FW (Kg)	4.30	1.99	0.42	12.08	11.66	46.19
FW/DW	3.56	0.31	2.80	5.82	3.02	8.80
DW(Kg)	1.21	0.54	0.14	3.05	2.91	44.84
PH(cm)	57.87	17.97	20.23	98.47	78.24	31.05
TBN	24.6	3.80	13	33	20	15.42
PW(cm)	158.18	21.05	75.32	223.21	147.89	14.21
LB(cm)	75.86	11.73	34.50	105.26	71.12	15.46
LL(cm)	3.24	0.27	2.54	3.98	1.44	8.31
LW(cm)	0.65	0.08	0.49	0.90	0.41	12.15
LL/LW	5.05	0.57	3.78	6.86	3.08	11.33