



Supplementary Material

Trends in Alcohol Intake and the Association between Socio-Demographic Factors and Volume of Alcohol Intake amongst Adult Male Drinkers in China

Ruiyi Liu ^{1,2,3,†}, Li Chen ^{1,2,3,†}, Fan Zhang ^{1,2,3}, Rui Zhu ^{1,2,3}, Xinjie Lin ⁴, Xuchen Meng ⁴, Huabing Li ^{1,2,3}, Xun Lei ^{1,2,3,*} and Yong Zhao ^{1,2,3,5*}

¹ School of Public Health and Management, Chongqing Medical University, Chongqing 400016, China; lry981118@foxmail.com (R.L.); nameclx@foxmail.com (L.C.); ava11@126.com (F.Z.); 2016111041@stu.cqmu.edu.cn (R.Z.); lhbcqmu@outlook.com (H.L.)

² Research Center for Medicine and Social Development, Chongqing Medical University, Chongqing 400016, China

³ The Innovation Center for Social Risk Governance in Health, Chongqing Medical University, Chongqing 400016, China

⁴ Clinical College, Chongqing Medical University, Chongqing 400016, China; linxinjie1@outlook.com (X.L.); angelitamxc@hotmail.com (X.M.)

⁵ Chongqing Key Laboratory of Child Nutrition and Health. Chongqing 400016, China;

† These authors contributed equally to this work.

* Correspondence: zhaoyong@cqmu.edu.cn (Y.Z.); leixun521@163.com (X.L.); Tel.: +86-23-6848-5008 (Y.Z.); +86-23-6848-5031 (X.L.)

Table S1. Trends in prevalence of different types of alcohol consumption among male drinkers by survey year.

Variable			2004	2006	2009	2011	Pearson Chi-Square	p-Value
Beer consumption	Non-beer-eater	N	1070	952	908	1121	79.81	0.000
		%	44.6	40.1	35.1	34.1		
	Beer-eater	N	1327	1423	1680	2167		
		%	55.4	59.9	64.9	65.9		
Wine consumption	Non-wine-eater	N	2172	2220	2292	2695	198.18	0.000
		%	90.6	93.5	88.6	82.0		
	Wine-eater	N	225	155	296	593		
		%	9.4	6.5	11.4	18.0		
Liqueur consumption	Non-liqueur-eater	N	413	501	594	809	47.30	0.000
		%	17.2	21.1	23.0	24.6		
	Liqueur-eater	N	1984	1874	1994	2479		
		%	82.8	78.9	77.0	75.4		

Table S2. Time trends in the prevalence of alcohol consumption (beer, wine, liqueur) amongst male drinkers by age groups, CHNS.

Age groups		Beer				Wine				Liqueur				
		2004	2006	2009	2011	2004	2006	2009	2011	2004	2006	2009	2011	
18-35	Non drinker	N	111	69	43	42	485	435	404	428	166	178	202	223
		%	20.4	14.8	9.3	8.1	89.0	93.5	87.1	82.1	30.5	38.3	43.5	42.8
36-50	drinker	N	434	396	421	479	60	30	60	93	379	287	262	298
		%	79.6	85.2	90.7	91.9	11.0	6.5	12.9	17.9	69.5	61.7	56.5	57.2
51-65	Non drinker	N	360	289	244	278	853	868	856	984	141	175	215	258
		%	38.8	31.6	25.4	23.5	91.8	95.0	89.3	83.2	15.2	19.1	22.4	21.8
≥66	drinker	N	569	625	715	905	76	46	103	199	788	739	744	925
		%	61.2	68.4	74.6	76.5	8.2	5.0	10.7	16.8	84.8	80.9	77.6	78.2
18-35	Non drinker	N	422	425	419	560	622	704	777	980	83	119	130	252
		%	61.4	55.9	47.9	46.2	90.5	92.6	88.9	80.9	12.1	15.7	14.9	20.8
36-50	drinker	N	265	335	455	652	65	56	97	232	604	641	744	960
		%	38.6	44.1	52.1	53.8	9.5	7.4	11.1	19.1	87.9	84.3	85.1	79.2
51-65	Non drinker	N	177	169	202	241	212	213	255	303	23	29	47	76
		%	75.0	71.6	69.4	64.8	89.8	90.3	87.6	81.5	9.7	12.3	16.2	20.4
≥66	drinker	N	59	67	89	131	24	23	36	69	213	207	244	296
		%	25.0	28.4	30.6	35.2	10.2	9.7	12.4	18.5	90.3	87.7	83.8	79.6



© 2019 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).