


## Supplementary Materials: A Simplified Genomic Profiling Approach Predicts outcome in Metastatic Colorectal Cancer

Carlo Capalbo, Francesca Belardinilli, Domenico Raimondo, Edoardo Milanetti, Umberto Malapelle, Pasquale Pisapia, Valentina Magri, Alessandra Prete, Silvia Pecorari, Mariarosaria Colella, Anna Coppa, Caterina Bonfiglio, Arianna Nicolussi, Virginia Valentini, Alessandra Tessitore, Beatrice Cardinali, Marialaura Petroni, Paola Infante, Matteo Santoni, Marco Filetti, Valeria Colicchia, Paola Paci, Silvia Mezi, Flavia Longo, Enrico Cortesi, Paolo Marchetti, Giancarlo Troncone, Diana Bellavia, Gianluca Canettieri and Giuseppe Giannini



Groups	N° pz	Median PFS (months)	Comparisons	$p$ value
Overall population	38	7	AGG vs PRG	<0.001
AGG	17	5	AGG vs NMG	0.555
PRG	11	15	PRG vs NMG	0.006
NMG	10	6		

**Figure S1.** Kaplan–Meier plot showing the impact of PRG (green), AGG (red) and NMG (blue) group stratification on progression free survival (PFS) in the KRAS WT mCRCs.