

Scientific reports Supplementary Information Figure S1

Bloodmeal host identification with inferences to feeding habits of a fish-fed mosquito, *Aedes baisasi*

Takashi Miyake*, Natsuki Aihara, Ken Maeda, Chuya Shinzato, Ryo Koyanagi,
Hirozumi Kobayashi & Kazunori Yamahira

*corresponding author, tmiyascb@gifu-u.ac.jp

Supplementary information Figure S1. Pictures of bloodmeal-host species of the mosquito *Aedes baisasi* in the Ryukyu Archipelago. (a) *Moringua microchir*; (b) *Pisodonophis boro*; (c) *Uropterygius concolor*; (d) *Gymnothorax pictus*; (e) *Periophthalmus argentilineatus*; (f) *Trypauchenopsis intermedia*; (g) *Mugilogobius* sp. ‘Izumi-haze’; (h) *Bostrychus sinensis*; (i) *Myersina macrostoma*; (j) *Entomacrodus striatus*; (k) *Istiblennius edentulus*; (l) *Salarias luctuosus*; (m) *Salarias fasciatus*; (n) *Blenniella bilitonensis*; (o) *Rhinecanthus verrucosus*; photo credits: a–c and e–n (Ken Maeda); d and o (Katsunori Tachihara).

