

Supplemental Table 1: Plasma and serum analyte values missing at random (MAR) or identified as outliers by the ROUT method.

Analyte	Within 12h		Day 1		Day 4		Day 7		Day 14	
	MAR	Outliers	MAR	Outliers	MAR	Outliers	MAR	Outliers	MAR	Outliers
IL-6	9 (21%)	0 (0%)	6 (14%)	1 (2%)	4 (10%)	0 (0%)	4 (10%)	0 (0%)	6 (14%)	0 (0%)
IL-8	2 (5%)	2 (5%)	4 (10%)	0 (0%)	6 (14%)	0 (0%)	5 (12%)	0 (0%)	3 (7%)	0 (0%)
G-CSF	1 (2%)	5 (12%)	4 (10%)	0 (0%)	5 (12%)	0 (0%)	5 (12%)	0 (0%)	4 (10%)	0 (0%)
SDF-1	0 (0%)	0 (0%)	1 (2%)	0 (0%)	4 (10%)	0 (0%)	3 (7%)	0 (0%)	0 (0%)	0 (0%)
TNF- α	3 (7%)	0 (0%)	5 (12%)	0 (0%)	3 (7%)	0 (0%)	3 (7%)	0 (0%)	1 (2%)	0 (0%)
EPO	1 (2%)	0 (0%)	5 (12%)	0 (0%)	4 (10%)	0 (0%)	5 (12%)	0 (0%)	3 (7%)	1 (2%)
sTfR	1 (2%)	0 (0%)	NA	NA	NA	NA	NA	NA	1 (2%)	0 (0%)

IL-6: interleukin 6, IL-8: interleukin 8, G-CSF: granulocyte colony-stimulating factor, SDF-1, stromal-derived factor 1, TNF- α : tumor necrosis factor alpha, EPO: erythropoietin, sTfR: serum transferrin receptor. Data are presented as n (%).