

Supplementary Table 3. Kaplan-Meier Analysis of Other Imaging and Clinico-Pathologic Parameters for DFS

Parameters	Threshold Values	Median Survival, Month (95% CI)		<i>P</i>
		Above Threshold or Presence	Below Threshold or Absence	
Age	> 47	31.00 (24.30–47.83)	65.70 (11.03–66.47)	0.232
Sex		26.83 (21.10–46.97)		0.068
Cirrhosis		24.80 (13.73–43.80)	46.97 (25.13–66.47)	0.081
Size	> 6.5	9.27 (5.73–13.33)	66.47 (43.80–66.47)	< 0.001
Microscopic vascular invasion		11.03 (6.47–29.20)	47.83 (31.00–66.47)	0.006
Satellite nodule		3.67 (3.17–11.80)	37.70 (25.80–66.47)	0.000
Histologic grade		13.73 (10.53–46.97)	47.83 (29.20–66.47)	0.038
AFP	> 14	25.80 (11.03–46.97)	66.47 (26.83–66.47)	0.021
PIVKA-II	> 61	17.90 (11.03–26.00)	65.70 (34.87–66.47)	0.012
Average attenuation of tumor (arterial-phase)	≤ 84.2	34.87 (24.80–66.47)	23.27 (11.03–29.20)	0.001
Average attenuation of tumor (portal phase)	≤ 100.65		26.83 (14.47–43.80)	0.036
Rim arterial enhancement		3.17 (2.87–17.90)	37.70 (25.80–66.47)	0.010
Peritumoral parenchymal enhancement		13.60 (3.17–45.87)	43.80 (25.13–66.47)	0.031
Non-smooth tumor margin		26.83 (21.10–46.97)	65.70 (23.27–66.47)	0.121

AFP = alpha-fetoprotein, PIVKA-II = vitamin K absence or antagonist-II