

**Supplementary Table 4. Interobserver Agreement (ICC) for CTTA on Arterial-Phase Images**

Filter Scale Values	Mean	SD	Entropy	MPP	Skewness	Kurtosis
No filtration	0.99 (0.98–1.00)	0.94 (0.88–0.97)	0.96 (0.92–0.98)	0.99 (0.98–0.99)	0.94 (0.87–0.97)	0.94 (0.86–0.97)
2.0 (fine)	0.99 (0.99–1.00)	0.83 (0.63–0.92)	0.99 (0.99–1.00)	0.94 (0.87–0.97)	0.78 (0.51–0.90)	0.99 (0.99–1.00)
3.0 (medium)	0.90 (0.78–0.95)	0.92 (0.81–0.96)	0.68 (0.29–0.85)	0.79 (0.54–0.91)	0.99 (0.99–1.00)	0.97 (0.93–0.99)
4.0 (medium)	0.82 (0.59–0.92)	0.90 (0.78–0.95)	0.94 (0.86–0.97)	0.90 (0.77–0.95)	0.76 (0.46–0.89)	0.97 (0.92–0.98)
5.0 (medium)	0.94 (0.86–0.97)	0.93 (0.84–0.97)	0.99 (0.99–1.00)	0.86 (0.69–0.94)	0.99 (0.99–1.00)	0.88 (0.72–0.95)
6.0 (coarse)	0.90 (0.75–0.96)	0.98 (0.95–0.99)	0.97 (0.93–0.99)	0.90 (0.77–0.96)	0.83 (0.60–0.93)	0.91 (0.78–0.96)

Values of ICC (95% CI). Guide to interpretation of ICCs for agreement: values < 0.40 were considered poor; 0.40–0.59, fair; 0.60–0.74, good; and 0.75–1.00, excellent. CTTA = computed tomography texture analysis, ICC = intraclass correlation coefficient