

Supplementary Table 5. Interobserver Agreement (ICC) for CTTA on Portal-venous-Phase Images

Filter Scale Values	Mean	SD	Entropy	MPP	Skewness	Kurtosis
No filtration	0.85 (0.67–0.93)	0.85 (0.67–0.93)	0.74 (0.44–0.88)	0.89 (0.75–0.95)	0.73 (0.41–0.88)	0.85 (0.68–0.93)
2.0 (fine)	0.94 (0.87–0.97)	0.97 (0.92–0.98)	0.84 (0.65–0.93)	0.84 (0.65–0.93)	0.78 (0.53–0.90)	0.78 (0.51–0.90)
3.0 (medium)	0.69 (0.32–0.86)	0.80 (0.56–0.91)	0.80 (0.56–0.91)	0.72 (0.39–0.87)	0.86 (0.69–0.94)	0.92 (0.83–0.97)
4.0 (medium)	0.99 (0.97–0.99)	0.89 (0.76–0.95)	0.91 (0.79–0.96)	0.94 (0.87–0.97)	0.78 (0.52–0.90)	0.69 (0.32–0.86)
5.0 (medium)	0.99 (0.99–1.00)	0.99 (0.99–1.00)	0.96 (0.90–0.98)	0.99 (0.99–1.00)	0.82 (0.59–0.92)	0.94 (0.86–0.97)
6.0 (coarse)	0.99 (0.98–1.00)	0.98 (0.96–1.00)	0.90 (0.76–0.96)	0.99 (0.97–1.00)	0.73 (0.36–0.89)	0.94 (0.85–0.97)

Values of ICC (95% CI). Guide to interpretation of ICCs for agreement: values < 0.40 were considered poor; 0.40–0.59, fair; 0.60–0.74, good; and 0.75–1.00, excellent.