

Amyloid beta in nasal secretions may be a potential biomarker of Alzheimer's disease

Young Hyo Kim,^{1*} Sang-Myung Lee,^{2*} Sungbo Cho,³ Ju-Hee Kang,⁴ Yang-Ki Minn,⁵ Hyelim Park,¹ Seong Hye Choi⁶

(*These authors contributed equally to this work.)

¹Departments of Otorhinolaryngology-Head and Neck Surgery, Inha University School of Medicine, Incheon22332, South Korea; ²Department of Chemical Engineering, Kangwon National University, Chuncheon24341, South Korea; ³Department of Electronic Engineering, Gachon University, Seongnam13120, South Korea; ⁴Department of Pharmacology, Inha University School of Medicine, Incheon22212, South Korea; ⁵Department of Neurology, Hallym University Kangnam Sacred Heart Hospital, Hallym University College of Medicine, Seoul07441, South Korea; ⁶Department of Neurology, Inha University School of Medicine, Incheon22332, South Korea

Corresponding author: Seong Hye Choi MD, PhD

Department of Neurology, Inha University Hospital
27 Inhang-ro, Jung-gu, Incheon 22332, South Korea
Tel: 82 32 890 3860
FAX: 82 32 890 1140
E-mail: seonghye@inha.ac.kr

