

# Molecular quantification of *Plasmodium* parasite density from the blood retained in used RDTs

Ailie Robinson\*

London School of Hygiene and Tropical Medicine, Keppel Street, London, WC1E 7HT, United Kingdom.  
[ailie.robinson@lshtm.ac.uk](mailto:ailie.robinson@lshtm.ac.uk)

Annette O Busula

Affiliations for this study: Laboratory of Entomology, Wageningen University & Research, 6708 PB, Wageningen, The Netherlands, International Centre of Insect Physiology and Ecology, Nairobi, Kenya. Current affiliation: Department of Biological and Agricultural Sciences, Kaimosi Friends University College, a constituent college of Masinde Muliro University of Science and Technology, Kaimosi, Kenya.

[abusula@kafuco.ac.ke](mailto:abusula@kafuco.ac.ke)

Julian K. Muwanguzi

London School of Hygiene & Tropical Medicine, Keppel Street, London, WC1E 7HT, United Kingdom.  
[Julian.Muwanguzi@lshtm.ac.uk](mailto:Julian.Muwanguzi@lshtm.ac.uk)

Stephen J. Powers

Affiliations for this study: Computational and Analytical Sciences, Rothamsted Research, Harpenden, Hertfordshire, AL5 2JQ, United Kingdom. Current affiliation: Rylands Farm, East Lambrook Road, South Petherton, Somerset, TA13 5HP, United Kingdom.

[powerssj@gmail.com](mailto:powerssj@gmail.com)

Daniel K. Masiga

Animal Health, International Centre of Insect Physiology and Ecology, PO Box 30772-00100, Nairobi, Kenya.

[dmasiga@icipe.org](mailto:dmasiga@icipe.org)

Teun Bousema

London School of Hygiene & Tropical Medicine, Keppel Street, London, WC1E 7HT, United Kingdom, and Medical Microbiology, Radboud University Medical Centre, Geert Grooteplein 26-28, 6525 GA Nijmegen, The Netherlands.

[Teun.Bousema@radboudumc.nl](mailto:Teun.Bousema@radboudumc.nl)

Willem Takken,

Laboratory of Entomology, Wageningen University & Research, PO Box 16, 6700 AA, Wageningen, The Netherlands.

[willem.takken@wur.nl](mailto:willem.takken@wur.nl)

Jetske G de Boer

Affiliation for this study: Laboratory of Entomology, Wageningen University & Research, 6708 PB, Wageningen, The Netherlands. Current affiliation: Netherlands Institute of Ecology, 6708 PB, Wageningen, The Netherlands.

[J.deBoer@nioo.knaw.nl](mailto:J.deBoer@nioo.knaw.nl)

James G. Logan

London School of Hygiene & Tropical Medicine, Keppel Street, London, WC1E 7HT, United Kingdom.

[James.Logan@lshtm.ac.uk](mailto:James.Logan@lshtm.ac.uk)

Khalid B. Beshir

London School of Hygiene & Tropical Medicine, Keppel Street, London, WC1E 7HT, United Kingdom.

[Khalid.Beshir@lshtm.ac.uk](mailto:Khalid.Beshir@lshtm.ac.uk)


Colin J. Sutherland

London School of Hygiene & Tropical Medicine, Keppel Street, London, WC1E 7HT, United Kingdom.

[Colin.Sutherland@lshtm.ac.uk](mailto:Colin.Sutherland@lshtm.ac.uk)

\*Corresponding author

## Supplementary Information



Supplementary Figure S1. The samples that were used in this study were collected from five- to 12-year-old-children, who were participating in a malaria odour study. Samples for that study were collected over three timepoints; in the present study parasite density was estimated using paired RDT and DBS blood samples from all available samples. Antimalarial treatment with artemether-lumefantrine was administered to all *Plasmodium*-positive individuals according to manufacturer's instructions.

Supplementary Table S1. The result of the RDT test (Pos P.f. or Pos P.f./pan grouped together as 'positive') in comparison to *Plasmodium* parasite positivity as determined by duplex qPCR based on two templates, fragments of the nitrocellulose membrane within a used rapid diagnostic test (RDT<sub>DNA</sub>) or dried blood spots collected onto Whatman filter paper Number 3 (DBS<sub>DNA</sub>). Column (1) shows the result of the RDT, columns (2) and (4) show agreement of the duplex qPCR (positive/negative) with RDT result according to the two templates, column (3) shows agreement of DBS<sub>DNA</sub> template qPCR with RDT<sub>DNA</sub> template qPCR, per RDT outcome.

1	2	3	4
RDT result, n (%)	qPCR RDT <sub>DNA</sub> template agreement with RDT result n (%)	DBS <sub>DNA</sub> template agreement with RDT <sub>DNA</sub> template, per RDT result n (%)	qPCR DBS <sub>DNA</sub> template agreement with RDT result n (%)
Positive: 86 (60.99)	Positive, 48 (55.81) <sup>(A)</sup>	Positive, 39 (81.25)	Positive, 47 (54.65)
		Inconclusive, 1 (2.08) <sup>(B)</sup>	
		Negative, 8 (16.67)	
	Inconclusive, 9 (10.47)	Positive, 3 (33.33)	Inconclusive, 9 (10.47)
		Inconclusive, 1 (11.11)	
		Negative, 5 (55.56)	
	Negative, 29 (33.72)	Negative, 17 (58.62)	Negative, 30 (34.88)
		Inconclusive, 7 (24.14)	
		Positive, 5 (17.24)	
Negative: 52 (36.88)	Negative, 23 (44.23)	Positive, 3 (13.04) <sup>(C)</sup>	Negative, 35 (67.31)
		Inconclusive, 5 (21.74)	
		Negative, 15 (65.44)	
	Inconclusive, 13 (25.0)	Positive, 1 (7.69)	Inconclusive, 11 (21.15)
		Inconclusive, 3 (23.08)	
		Negative, 9 (69.23)	
	Positive, 16 (30.77) <sup>(D)</sup>	Positive, 2 (12.5) <sup>(E)</sup>	Positive, 6 (11.54)
		Inconclusive, 3 (18.75)	
		Negative, 11 (68.75)	
Not recorded: 3 (2.13)	Inconclusive, 1 (33.33)	Negative, 1 (100)	Negative, 2 (66.67)
	Negative, 2 (66.67)	Negative, 1 (50)	
		Inconclusive, 1 (50)	

<sup>(A)</sup> qPCR positive samples had detectable parasite DNA in each of two technical replicates.

<sup>(B)</sup> Inconclusive samples were those with one positive and one negative replicate.

<sup>(C)</sup> Samples were 6, 25.33, 187.50 p/μL

<sup>(D)</sup> Mean/median: 26.73/0.02 p/μL

<sup>(E)</sup> Samples were 12.47 and 8.76 p/μL