

Parameter	Time	CKD patients			Healthy volunteers			
		2 mg	4 mg	8 mg	1.2 mg	4 mg	8 mg	16 mg
Hb g/dl	19/24h*	0.62	1.32	1.23	-0.05	0.23	0.10	0.12
	44/48h*	0.02	0.42	0.57	-0.17	0.1	0.13	-0.12
	168h	-0.40	0.06	0.25				
	240h				-0.15	-0.08	0.10	-0.32
	672h	-0.58	0.30	0.48	0.10	0.12	0.22	0.07
Ret 0/00	19/24h*	-1.26	-0.84	0.05	0.01	0.08	0.02	-0.01
	44/48h*	-3.53	-0.67	-0.78	0.06	0.05	0.06	0.06
	168h	-1.40	0.96	2.90				
	240h				0.15	0.18	-0.06	0.44
	672h	-2.62	-0.30	-0.07	0.13	-0.22	-0.01	0.04
RetHb pg	19/24h*	-0.14	-2.04	0.07	-0.01	0.05	0.03	-0.02
	44/48h*	-1.10	-0.13	0.83	0.02	-0.02	0.05	0.03
	168h	0.5	-0.16	0.83				
	240h				-0.03	-0.05	-0.03	0.05
	672h	-0.77	0.43	0.10	-0.07	-0.03	-0.13	-0.02

\*In CKD patients, slow response parameters were assessed at 19 and 44 hours, in healthy volunteers at 24 and 48 hours