

Table S2. Solicited reactions and associated intensity scales

Reaction	Intensity Scale
Local	
Injection-site tenderness	Grade 1: Minor reaction when injection site is touched Grade 2: Cries or protests when injection site is touched Grade 3: Cries when injected limb is moved, or the movement of the injected limb is reduced
Injection-site erythema and injection-site swelling	Grade 1: > 0 to < 25 mm Grade 2: \geq 25 to < 50 mm Grade 3: \geq 50 mm
Systemic	
Fever	Grade 1: \geq 100.4°F to \leq 101.3°F Grade 2: > 101.3°F to \leq 103.1°F Grade 3: > 103.1°F
Vomiting	Grade 1: 1 episode per 24 hours Grade 2: 2–5 episodes per 24 hours Grade 3: \geq 6 episodes per 24 hours or requiring parenteral hydration

Abnormal crying

Grade 1: < 1 hour

Grade 2: 1–3 hours

Grade 3: > 3 hours

Drowsiness

Grade 1: Sleepier than usual or less interested in surroundings

Grade 2: Not interested in surroundings or did not wake up for a feed/meal

Grade 3: Sleeping most of the time or difficult to wake up

Loss of appetite

Grade 1: Eating less than normal

Grade 2: Missed 1 or 2 feeds/meals completely

Grade 3: Refuses ≥ 3 feeds/meals or refuses most feeds/meals

Irritability

Grade 1: Easily consolable

Grade 2: Requiring increased attention

Grade 3: Inconsolable
