

**Table S7. Grade 3 solicited reactions within 7 days**

<b>Grade 3 solicited reaction</b>	<b>Half dose of IIV4</b>		<b>Full dose of IIV4</b>	
	<b>n/N</b>	<b>% (95% CI)</b>	<b>n/N</b>	<b>% (95% CI)</b>
Any	67/909	7.4 (5.8, 9.3)	79/941	8.4 (6.7, 10.4)
Injection site				
Tenderness	16/909	1.8 (1.0, 2.8)	13/939	1.4 (0.7, 2.4)
Tenderness	15/909	1.7 (0.9, 2.7)	11/939	1.2 (0.6, 2.1)
Redness	0/909	0.0 (0.0, 0.4)	2/938	0.2 (0.0, 0.8)
Swelling	1/908	0.1 (0.0, 0.6)	0/937	0.0 (0.0, 0.4)
Systemic				
Fever	58/909	6.4 (4.9, 8.2)	67/941	7.1 (5.6, 9.0)
Fever	5/893	0.6 (0.2, 1.3)	11/930	1.2 (0.6, 2.1)
Vomiting	4/908	0.4 (0.1, 1.1)	5/941	0.5 (0.2, 1.2)
Abnormal crying	28/908	3.1 (2.1, 4.4)	24/941	2.6 (1.6, 3.8)
Drowsiness	19/908	2.1 (1.3, 3.2)	15/940	1.6 (0.9, 2.6)
Loss of appetite	13/908	1.4 (0.8, 2.4)	21/940	2.2 (1.4, 3.4)
Irritability	33/908	3.6 (2.5, 5.1)	38/940	4.0 (2.9, 5.5)

Safety was assessed in all vaccinated participants according to the vaccine they actually received.

Abbreviations: AE, adverse event; SAE, serious adverse event.