

## Supplementary Online Content

Meghji Z, Nguyen A, Fatima B, et al. Survival differences in women and men after septal myectomy for obstructive hypertrophic cardiomyopathy. *JAMA Cardiol*. Published online February 27, 2019. doi:10.1001/jamacardio.2019.0084

**eTable.** Baseline Characteristics over the Study Period

**eFigure.** Temporal trend in baseline characteristics by sex. Curve estimates trend over time stratified by sex using a locally weighted scatterplot smooth (loess).

This supplementary material has been provided by the authors to give readers additional information about their work.

**eTable. Baseline Characteristics over the Study Period**

<b>Variable</b>	<b>N</b>	<b>Women</b>	<b>Men</b>
<b>Age</b>			
1960-79	49	47.5 (37.9, 63.9)	43.0 (30.9, 53.2)
1980-89	76	58.1 (36.4, 67.1)	48.9 (37.0, 61.9)
1990-99	276	59.2 (41.6, 68.6)	49.7 (38.4, 61.5)
2000-09	985	56.9 (45.8, 68.8)	52.0 (43.4, 62.0)
2010-16	1,129	61.1 (49.8, 68.1)	54.8 (44.9, 63.7)
<b>NYHA class III or IV</b>			
1960-79	49	16 (72.7%)	20 (74.1%)
1980-89	76	33 (91.7%)	36 (90.0%)
1990-99	276	132 (93.6%)	112 (88.9%)
2000-09	985	364 (87.9%)	474 (83.0%)
2010-16	1,129	478 (93.0%)	527 (85.7%)
<b>Resting LVOT gradient</b>			
1960-79	-	-	-
1980-89	55	64.0 (36.0, 100.0)	64.0 (36.0, 94.0)
1990-99	225	76.0 (46.0, 102.0)	61.0 (36.0, 92.0)
2000-09	965	67.0 (36.0, 100.0)	49.0 (21.0, 81.0)
2010-16	1,089	64.0 (29.0, 92.0)	49.0 (19.0, 77.0)
<b>MR grade (moderate to severe)</b>			
1960-79	-	-	-
1980-89	68	23 (69.7%)	19 (54.3%)
1990-99	261	79 (57.7%)	61 (49.2%)
2000-09	980	236 (57.4%)	235 (41.3%)
2010-16	1,129	267 (51.9%)	264 (42.9%)

**eFigure.** Temporal trend in baseline characteristics by sex. Curve estimates trend over time stratified by sex using a locally weighted scatterplot smooth (loess).

