Supplementary Online Content

Berlinberg EJ, Gonzales JA, Doan T, Acharya NR. Association between noninfectious uveitis and psychological stress. *JAMA Ophthalmol*. Published online December 6, 2018. doi:10.1001/jamaophthalmol.2018.5893

eTable. Demographic and Clinical Characteristics of Uveitis Group

This supplementary material has been provided by the authors to give readers additional information about their work.

eTable. Demographic and Clinical Characteristics of Uveitis Group

Variable	Recently Active	Controlled Uveitis	<i>P</i> -value
	Uveitis N (%)	N (%)	
Female	24 (60%)	26 (65%)	.82ª
Mean Age (SD)	43.4 (18.6)	45.4 (16.1)	.60 ^b
Race/Ethnicity			.68ª
White	21 (52 %)	23 (58%)	
Black	4 (10%)	3 (8%)	
Asian	7 (18%)	10 (25%)	
Hispanic	3 (8%)	3 (8%)	
Other	1 (2%)	0 (0%)	
Multiracial	4 (10%)	1 (2%)	
Current smoker	2 (5%)	2 (5%)	.85ª
Median Household Income in Dollars	86,661.50	78,301.00	.90°
(IQR)	(56,646.50 –	(68,869.00 –	
	129,108.00)	119,281.00)	
Marital Status	•		.29ª
Single	18 (45%)	13 (32%)	
Married	21 (52%)	22 (55%)	
Divorced	0 (0%)	3 (8%)	
Widowed	1 (2%)	1 (2%)	
Separated	0 (0%)	1 (2%)	
Living Situation	, ,	· · · · · · · · · · · · · · · · · · ·	.78ª
Alone	4 (10%)	7 (18%)	
With Family	29 (72%)	26 (65%)	
With Friends	2 (5%)	3 (8%)	
With Roommates	5 (12%)	4 (10%)	
Education		,	.27ª
Less than High School	1 (2%)	0 (0%)	
High School	3 (8%)	3 (8%)	
Some College	13 (32%)	6 (15%)	
Bachelor's Degree	12 (30%)	15 (38%)	
Professional or Graduate Degree	11 (28%)	16 (15%)	
Mean Drinks per Week (SD)	2 (3)	2 (3)	.89 ^b
Have Dependents	11 (28%)	12 (30%)	>.99 ^a
Comorbid Conditions	\/	\/	
Heart disease	3 (8%)	1 (2%)	.62ª
Cancer	5 (12%)	4 (10%)	>.99 ^a
Depression	12 (30%)	4 (10%)	.05ª
Anxiety	9 (22%)	6 (15%)	.57ª
Hypertension	7 (18%)	3 (8%)	.31ª
Diabetes	3 (8%)	3 (8%)	>.99ª
Systemic Autoimmune (Other than Diabetes)	16 (40%)	20 (50%)	.50ª

Median Number of All Other Comorbid Conditions (Excluding Systemic Autoimmune) (IQR) 0 (0 - 1) 0.5 (0 - 2) Work Status 4 (10%) 2 (5%) Full Time 21 (52%) 24 (60%) Part Time 5 (12%) 4 (10%) Retired 7 (18%) 7 (18%) Homemaker 0 (0%) 2 (5%) Other 3 (8%) 1 (2%) VAS Pain (0-10) mean (SD) 20.1 (26.1) 12.4 (20.2) Median converted logMAR Vision in the better seeing eye (IQR) 0.14 (0 - 0.39) 0 (0 - 0.1) Anatomical Location 23 (58%) 18 (45%) Intermediate 2 (2%) 5 (12%) Anterior + Intermediate 3 (8%) 7 (18%)	.08°
Conditions (Excluding Systemic Autoimmune) (IQR) 4 (10%) 2 (5%) Work Status 21 (52%) 24 (60%) Full Time 21 (52%) 24 (60%) Part Time 5 (12%) 4 (10%) Retired 7 (18%) 7 (18%) Homemaker 0 (0%) 2 (5%) Other 3 (8%) 1 (2%) VAS Pain (0-10) mean (SD) 20.1 (26.1) 12.4 (20.2) Median converted logMAR Vision in the better seeing eye (IQR) 0.14 (0 - 0.39) 0 (0 - 0.1) Anatomical Location 23 (58%) 18 (45%) Intermediate 2 (2%) 5 (12%)	.66ª
Work Status 4 (10%) 2 (5%) Full Time 21 (52%) 24 (60%) Part Time 5 (12%) 4 (10%) Retired 7 (18%) 7 (18%) Homemaker 0 (0%) 2 (5%) Other 3 (8%) 1 (2%) VAS Pain (0-10) mean (SD) 20.1 (26.1) 12.4 (20.2) Median converted logMAR Vision in the better seeing eye (IQR) 0.14 (0 - 0.39) 0 (0 - 0.1) Anatomical Location 23 (58%) 18 (45%) Intermediate 2 (2%) 5 (12%)	
Full Time 21 (52%) 24 (60%) Part Time 5 (12%) 4 (10%) Retired 7 (18%) 7 (18%) Homemaker 0 (0%) 2 (5%) Other 3 (8%) 1 (2%) VAS Pain (0-10) mean (SD) 20.1 (26.1) 12.4 (20.2) Median converted logMAR Vision in the better seeing eye (IQR) 0.14 (0 - 0.39) 0 (0 - 0.1) Anatomical Location 23 (58%) 18 (45%) Intermediate 2 (2%) 5 (12%)	14 ^b
Full Time 21 (52%) 24 (60%) Part Time 5 (12%) 4 (10%) Retired 7 (18%) 7 (18%) Homemaker 0 (0%) 2 (5%) Other 3 (8%) 1 (2%) VAS Pain (0-10) mean (SD) 20.1 (26.1) 12.4 (20.2) Median converted logMAR Vision in the better seeing eye (IQR) 0.14 (0 - 0.39) 0 (0 - 0.1) Anatomical Location 23 (58%) 18 (45%) Intermediate 2 (2%) 5 (12%)	14b
Part Time 5 (12%) 4 (10%) Retired 7 (18%) 7 (18%) Homemaker 0 (0%) 2 (5%) Other 3 (8%) 1 (2%) VAS Pain (0-10) mean (SD) 20.1 (26.1) 12.4 (20.2) Median converted logMAR Vision in the better seeing eye (IQR) 0.14 (0 - 0.39) 0 (0 - 0.1) Anatomical Location 23 (58%) 18 (45%) Intermediate 2 (2%) 5 (12%)	14b
Retired 7 (18%) 7 (18%) Homemaker 0 (0%) 2 (5%) Other 3 (8%) 1 (2%) VAS Pain (0-10) mean (SD) 20.1 (26.1) 12.4 (20.2) Median converted logMAR Vision in the better seeing eye (IQR) 0.14 (0 - 0.39) 0 (0 - 0.1) Anatomical Location 23 (58%) 18 (45%) Intermediate 2 (2%) 5 (12%)	14 ^b
Homemaker 0 (0%) 2 (5%) Other 3 (8%) 1 (2%) VAS Pain (0-10) mean (SD) 20.1 (26.1) 12.4 (20.2) Median converted logMAR Vision in the better seeing eye (IQR) 0.14 (0 - 0.39) 0 (0 - 0.1) Anatomical Location 23 (58%) 18 (45%) Intermediate 2 (2%) 5 (12%)	14 ^b
Other 3 (8%) 1 (2%) VAS Pain (0-10) mean (SD) 20.1 (26.1) 12.4 (20.2) Median converted logMAR Vision in the better seeing eye (IQR) 0.14 (0 - 0.39) 0 (0 - 0.1) Anatomical Location 23 (58%) 18 (45%) Intermediate 2 (2%) 5 (12%)	14 ^b
VAS Pain (0-10) mean (SD) 20.1 (26.1) 12.4 (20.2) Median converted logMAR Vision in the better seeing eye (IQR) 0.14 (0 - 0.39) 0 (0 - 0.1) Anatomical Location 23 (58%) 18 (45%) Intermediate 2 (2%) 5 (12%)	14 ^b
Median converted logMAR Vision in the better seeing eye (IQR) 0.14 (0 - 0.39) 0 (0 - 0.1) Anatomical Location 23 (58%) 18 (45%) Intermediate 2 (2%) 5 (12%)	T
better seeing eye (IQR) Anatomical Location Anterior 23 (58%) 18 (45%) Intermediate 2 (2%) 5 (12%)	.003°
Anatomical Location 23 (58%) 18 (45%) Intermediate 2 (2%) 5 (12%)	
Intermediate 2 (2%) 5 (12%)	.16ª
Intermediate 2 (2%) 5 (12%)	
Posterior 6 (15%) 2 (5%)	
Panuveitis 7 (18%) 8 (20%)	
Bilateral Disease 35 (88%) 28 (70%)	.10 ^a
Uveitis Diagnosis	.09ª
Undifferentiated 20 (50%) 11 (28%)	
Retinal vasculitis 2 (5%) 2 (5%)	
Serpiginous choroiditis 0 (0%) 1 (2%)	
Multifocal choroiditis with panuveitis 2 (5%) 2 (5%)	
Birdshot retinochoroidopathy 0 (0%) 2 (5%)	
Pars planitis 3 (8%) 1 (2%)	
Behçet's disease 0 (0%) 2 (5%)	
Sarcoidosis 3 (8%) 4 (10%)	
Multiple sclerosis 0 (0%) 2 (5%)	
Juvenile idiopathic arthritis 1 (2%) 1 (2%)	
Seronegative spondyloarthropathy 2 (5%) 8 (20%)	
Inflammatory bowel disease 0 (0%) 1 (2%)	
HLA-B27 positive, no reported joint 0 (0%) 3 (8%) involvement	
Vogt-Koyanagi-Harada disease 0 (0%) 1 (2%)	
Other 4 (10%) 2 (5%)	
Median years duration of disease (IQR) 5.5 (1 - 12) 6 (2 - 14)	.30°
Diagnosed Systemic Disease 16 (40%) 23 (58%)	.18 ^a
Any corticosteroid use 33 (82%) 17 (42%)	<.001 ^a
Current Immunomodulatory Therapy 18 (45%) 20 (50%)	.82ª
Number of eye medications (IQR) 1.5 (1 - 3.5) 1 (0 - 3)	.04
Number of previous injections (IQR) 0 (0 - 1.5) 0 (0 - 0)	.17°

a. Fisher's exact test

b. Student's t-testc. Mann-Whitney U test