

Supplementary Online Content

Howard R, Fry B, Gunaseelan V, et al. Association of opioid prescribing with opioid consumption after surgery in Michigan. *JAMA Surg*. Published online November 7, 2018. doi:10.1001/jamasurg.2018.4234

eTable. Leftover Opioids by Procedure

This supplementary material has been provided by the authors to give readers additional information about their work.

eTable. Leftover Opioids by Procedure

	Leftover Pills (Median (IQR))	Total Leftover Pills	Leftover OME (Median (IQR))	Total Leftover OME
Laparoscopic cholecystectomy	18 (5-28)	10636	100 (35-150)	61529
Laparoscopic appendectomy	20 (5-29)	4154	100 (37-150)	24858
Inguinal/femoral hernia repair (open/lap)	20 (5-29)	11943	100 (25-150)	70170
Open incisional hernia repair	15 (0-30)	2559	90 (0-150)	16209
Laparoscopic Colectomy	20 (10-30)	2251	114 (50-200)	14109
Open colectomy	20 (0-30)	1892	111 (0-163)	12626
Ileostomy/Colostomy	10 (0-27)	859	75 (0-200)	6388
Small Bowel Resection/Enterolysis	15 (0-29)	464	75 (0-150)	2628
Thyroidectomy	24 (13-30)	877	128 (75-161)	5668
Vaginal Hysterectomy	19 (5-27)	1870	100 (23-150)	13094
Laparoscopic/Robotic Hysterectomy	20 (5-30)	3804	120 (40-150)	22457
Abdominal Hysterectomy	15 (0-30)	1383	75 (0-150)	8713
TOTAL:		42692		258449