

S4 Table. Correlation among inflammatory biomarkers, stratified by HIV serostatus

HIV-infected men (N=452)										
	sCD163	sCD14	CCL2	IL-6	ICAM-1	Fibrinogen	D-dimer	CRP	sTNF- α R1	sTNF- α R2
sCD163	1.00									
sCD14	0.12	1.00								
CCL2	0.15	0.19	1.00							
IL-6	0.10	0.14	0.05	1.00						
ICAM-1	0.46	0.21	0.16	0.15	1.00					
Fibrinogen	0.06	0.12	0.10	0.12	0.06	1.00				
D-dimer	0.13	0.20	0.05	0.21	0.12	0.11	1.00			
CRP	0.04	0.27	0.16	0.28	0.15	0.21	0.45	1.00		
sTNF-αR1	0.14	0.21	0.25	0.10	0.12	0.06	0.16	0.09	1.00	
sTNF-αR2	0.42	0.38	0.33	0.24	0.43	0.13	0.33	0.35	0.71	1.00
HIV-uninfected men (N=276)										
	sCD163	sCD14	CCL2	IL-6	ICAM-1	Fibrinogen	D-dimer	CRP	sTNF- α R1	sTNF- α R2
sCD163	1.00									
sCD14	0.08	1.00								
CCL2	0.06	0.07	1.00							
IL-6	0.15	0.23	0.30	1.00						
ICAM-1	0.34	0.06	0.05	0.11	1.00					
Fibrinogen	0.18	0.21	0.15	0.26	0.16	1.00				
D-dimer	0.06	0.16	0.23	0.44	0.06	0.26	1.00			
CRP	0.18	0.28	0.22	0.50	0.19	0.52	0.47	1.00		
sTNF-αR1	0.21	0.11	0.29	0.28	0.17	0.17	0.27	0.20	1.00	
sTNF-αR2	0.35	0.21	0.22	0.21	0.34	0.17	0.09	0.25	0.61	1.00

Pearson correlation coefficient included in tables with 1.0 indicating a perfect positive correlation. Abbreviations: sCD163, cluster of differentiation 163; sCD14, cluster of differentiation 14; CCL2, chemokine (C-C motif) ligand 2; IL-6, interleukin-6; ICAM-1, intercellular cell adhesion molecule-1; CRP, C reactive protein; sTNF- α R1, tumor necrosis factor-alpha receptor 1; sTNF- α R2, tumor necrosis factor-alpha receptor 2.