

Optimal blood pressure after reperfusion therapy in patients with acute ischemic stroke

Kang-Ho Choi, MD, PhD^{1,2*†}, Jae-Myung Kim, MD^{1†}, Ja-Hae Kim, MD, PhD^{3,4*}, Joon-Tae Kim, MD, PhD¹, Man-Seok Park, MD, PhD¹, Seong-Min Choi, MD, PhD¹, Seung-Han Lee, MD, PhD¹, Byeong C. Kim, MD, PhD¹, Myeong-Kyu Kim, MD, PhD¹, Ki-Hyun Cho, MD, PhD¹

¹Department of Neurology, Chonnam National University Hospital, Gwangju, Korea

²Department of Neurology, Chonnam National University Hwasun Hospital, Hwasun, Korea

³Department of Nuclear Medicine, Chonnam National University Hospital, Gwangju, Korea

⁴Molecular Imaging Center, Chonnam National University Hwasun Hospital, Hwasun, Korea

***Authors for correspondence and requests for reprints:**

Kang-Ho Choi, MD, PhD

Department of Neurology, Chonnam National University Hwasun Hospital 322

Seoyang-Ro, Hwasun, Chonnam, Republic of Korea

Tel.: +82-62-220-6137, fax: +82-62-228-3461

E-mail: ckhchoikang@hanmail.net, ckhchoikang@chonnam.ac.kr

Ja-Hae Kim, MD, PhD

Department of Nuclear Medicine, Chonnam National University Hospital

8 Hakdong, Donggu, Gwangju, Republic of Korea


Tel.: +82-62-220-6155

E-mail: jhbt0607@hanmail.net

[†]These authors contributed equally to this work.

Supplementary figure

eFigure 1. (A) Distribution of modified Rankin Scale scores according to systolic blood pressure. (B) Optimal cutoff values for systolic and diastolic blood pressures for favorable outcomes.



ROC, receiver operating characteristic; AUC, area under the curve.