Table Supplemental Digital Content 1.

Frequency of All Pain Quality Descriptors Selected by the Total Sample of Women with Generalized Vulvodynia and Provoked Vestibulodynia (n = 60)

Neuropathic (n = 28 words)		Nociceptive (n = 26 words)	Other (n = 24 words)	
Burning	52 (87%)	Heavy	36 (60%)	Punishing	45 (75%)
Itchy	43 (72%)	Pressing	36 (60%)	Cruel	38 (63%)
Hot	41 (68%)	Crushing	34 (57%)	Annoying	38 (63%)
Stabbing	41 (68%)	Sharp	34 (57%)	Dreadful	37 (62%)
Aching	39 (65%)	Hurting	33 (55%)	Killing	36 (60%)
Tight	35 (58%)	Squeezing	31 (52%)	Terrifying	29 (48%)
Stinging	34 (57%)	Tearing	28 (47%)	Vicious	29 (48%)
Tingling	30 (50%)	Lacerating	22 (37%)	Troublesome	27 (45%)
Boring	28 (47%)	Cramping	20 (33%)	Torturing	27 (45%)
Pricking	25 (42%)	Piercing	20 (33%)	Unbearable	26 (43%)
Lancinating	23 (38%)	Cutting	19 (32%)	Miserable	22 (36%)
Searing	22 (37%)	Pinching	18 (30%)	Sickening	20 (33%)
Cold	16 (27%)	Pounding	18 (30%)	Frightening	18 (30%)
Cool	12 (20%)	Beating	13 (22%)	Grueling	12 (20%)
Freezing	7 (12%)	Pulsing	11 (18%)	Fearful	9 (15%)
Scalding	6 (10%)	Splitting	11 (18%)	Agonizing	9 (15%)
Penetrating	4 (7%)	Tender	11 (18%)	Tiring	8 (13%)
Shooting	4 (7%)	Dull	10 (17%)	Suffocating	8 (13%)
Flickering	3 (5%)	Sore	10 (17%)	Exhausting	7 (11%)
Drilling	2 (3%)	Gnawing	8 (13%)	Intense	7 (11%)
Quivering	2 (3%)	Throbbing	8 (13%)	Blinding	6 (10%)
Radiating	2 (3%)	Wrenching	4 (7%)	Nagging	3 (5%)
Smarting	2 (3%)	Pulling	3 (5%)	Nauseating	2 (3%)
Spreading	2 (3%)	Tugging	3 (5%)	Wretched	1 (2%)
Drawing	1 (2%)	Rasping	1 (2%)		
Flashing	1 (2%)	Taut	1 (2%)		
Jumping	1 (2%)				
Numb	1 (2%)				