

Table S3. Primers and probes for RT-qPCR used in this study

ZIKV-anti	GGTCAGCGTCCTCTCTAATAAACG
ZIKV-sense	GCACCCTAGTGTCCACTTTTTCC
ZIKV-Probe	FAM-AGCCATGACCGACACCACACCGT-BQ1
INF α -anti	GCCTCGCCCTTTGCTTTACT
INF α -sense	CTGTGGGTCTCAGGGAGATCA
INF β -anti	ATGACCAACAAGTGTCTCCTCC
INF β -sense	GGAATCCAAGCAAGTTGTAGCTC
IFN γ -anti	TCGGTAACTGACTTGAATGTCCA
IFN γ -sense	TCGCTTCCCTGTTTTAGCTGC
ISG15-anti	GAGAGGCAGCGAACTCATCT
ISG15-sense	CTTCAGCTCTGACACCGACA
ISG56-anti	TTGATGACGATGAAATGCCTGA
ISG56-sense	CAGGTCACCAGACTCCTCAC
IFITM3-anti	CATCCTCATGACCATTCTGC
IFITM3-sense	TCAGTGATGCCTCCTGATCT
Mx1-anti	GTTTCCGAAGTGGACATCGCA
Mx1-sense	CTGCACAGGTTGTTCTCAGC