

Effects of rikkunshito on renal fibrosis and inflammation in angiotensin II-infused mice

Kengo Azushima^{1,2†*}, Kazushi Uneda^{1†}, Hiromichi Wakui^{1*}, Kohji Ohki¹, Kotaro Haruhara¹,
Ryu Kobayashi¹, Sona Haku¹, Sho Kinguchi¹, Takahiro Yamaji¹, Shintaro Minegishi¹,
Tomoaki Ishigami¹, Akio Yamashita³ and Kouichi Tamura¹

† These authors contributed equally to this work.

¹ Department of Medical Science and Cardiorenal Medicine, Yokohama City University
Graduate School of Medicine, Yokohama, Japan

² Cardiovascular and Metabolic Disorders Program, Duke-NUS Medical School, Singapore

³ Department of Molecular Biology, Yokohama City University Graduate School of Medicine,
Yokohama, Japan

* Corresponding authors: Kengo Azushima, M.D., Ph.D. and Hiromichi Wakui, M.D., Ph.D.

Department of Medical Science and Cardiorenal Medicine

Yokohama City University Graduate School of Medicine

3-9 Fukuura, Kanazawa-ku, Yokohama, Japan 236-0004.

Tel.: 81-45-787-2635; Fax: 81-45-701-3738


E-mail: azushima@yokohama-cu.ac.jp or hiro1234@yokohama-cu.ac.jp

Full unedited gel for Figure 5D

GLP-1



β -actin



C, Control group; A, Ang II group; R, Ang II + RKT group.