

**Supplement Table 1** Demographic, socio-economic, and environmental characteristics of children hospitalised with severe pneumonia, stratified by age group and outcome

	<u>Children &lt;2 months</u>				<u>Children 2–59 months</u>			
	Survived n=709	Died n=40	CFR 5.3	p-value	Survived n=4107	Died n=198	CFR 4.6	p-value
<b>Age group</b>								
<b>Less than 1 month</b>	286 (40.3)	17 (42.5)	5.6	0.860				<0.001
<b>1 month</b>	423 (59.7)	23 (57.5)	5.2					
<b>2-5 months</b>					1011 (25.2)	76 (38.4)	7.0	
<b>6-11months</b>					1064 (25.9)	50 (25.3)	4.5	
<b>12-35 months</b>					1677 (40.8)	59 (29.8)	3.4	
<b>36-59 months</b>					355 (8.6)	13 (6.7)	3.5	
<b>Gender</b>								
<b>Female</b>	285 (40.2)	17 (42.5)	5.6	0.900	1786 (43.5)	90(45.5)	4.8	0.490
<b>Male</b>	424 (59.8)	23 (57.5)	5.1		2321 (56.5)	108 (54.5)	4.5	
<b>Hospital admitted</b>								
<b>BPH</b>	215 (30.3)	3 (7.5)	1.4	<0.001	795 (19.4)	26 (13.1)	3.2	<0.001
<b>EVRMC</b>	380 (53.6)	32 (80.0)	7.8		2429 (59.1)	148 (74.7)	5.7	
<b>ONP</b>	88 (12.4)	4 (10.0)	4.3		712 (17.3)	15 (7.6)	2.1	
<b>RITM</b>	26 (3.7)	1 (2.5)	3.7		171 (4.2)	9 (4.6)	5.0	
<b>Congenital abnormalities</b>								
<b>Yes</b>	8 (1.1)	3 (2.5)	27.3	0.010	33 (0.8)	9 (4.6)	6.2	<0.001
<b>No</b>	691 (98.9)	37 (92.5)			4035 (99.2)	187 (95.4)		
<b>Missing</b>	10	0			39	2		
<b>Height(h)-for-age</b>								
<b>-2SD ≤ h≤2SD</b>	251 (59.1)	5 (45.5)	2.0	0.350	1213 (53.9)	33 (48.5)	2.6	0.320
<b>-3SD ≤ h&lt;-2SD</b>	63 (14.8)	1 (9.1)	1.6		397 (17.6)	10(14.7)	2.5	
<b>h&lt;-3SD</b>	111 (26.1)	5 (45.5)	4.3		642 (28.5)	25(36.8)	3.7	
<b>Missing</b>	284	29			1855	130		
<b>Weight(w)-for-age</b>								
<b>-2SD ≤ w≤2SD</b>	511 (78.7)	15 (44.1)	2.9	<0.001	2424 (61.2)	82 (43.3)	3.3	<0.001
<b>-3SD ≤ w&lt;-2SD</b>	86 (13.3)	5 (14.7)	5.5		764 (19.3)	36 (18.6)	4.5	
<b>w&lt;-3SD</b>	52 (8.0)	14 (41.2)	21.2		776 (19.6)	76 (39.2)	8.9	

Missing	61 8 [6–9]	6 3.5 [1–8.2]	<0.001	143 4 [3–6]	6 1 [1–3]	<0.001		
<b>Duration of hospitalisation</b>								
<b>Environmental factors*</b>								
	n=184	n=4		n=904	n=23			
<b>SES</b>								
<b>≤30</b>	83 (46.1)	3 (75.0)	3.5	0.560	504 (55.8)	15 (65.2)	2.9	0.490
<b>&gt;30</b>	97 (53.9)	1 (25.0)	1.0		400 (44.2)	8 (34.4)	2.0	
<b>Missing</b>	5	1		0	0	0		
<b>Past experience of child loss</b>								
<b>Yes</b>	15 (11.4)	0 (0.0)	0.0	1.000	71 (10.2)	7 (33.3)	9.0	0.003
<b>No</b>	117 (88.6)	2 (5.0)	1.7		626 (89.8)	14 (66.7)	2.2	
<b>Missing</b>	53	3		207	2			
<b>Number of siblings</b>								
<b>&lt;2</b>	55 (29.7)	2 (50.0)	3.5	0.420	199 (22.1)	3 (13.5)	1.5	0.050
<b>2–3</b>	78 (42.2)	2 (50.0)	2.5		441 (48.9)	8 (34.8)	1.8	
<b>≥4</b>	52 (28.1)	0 (0.0)	0.0		262 (29.0)	12 (52.2)	4.4	
<b>Missing</b>	0	1		2	0			
<b>Types of cooking fuel used</b>								
<b>Electricity, petroleum gas, and Kerosene</b>	26 (14.1)	0	0.0	0.100	121 (12.8)	2 (8.7)	1.7	0.780
<b>Charcoal, and Wood</b>	158 (85.9)	4 (100.0)	2.5		775 (86.5)	21 (91.3)	2.6	
<b>Missing</b>	1	1		8	23			

NOTE: n (%), number and percent of child unless stated otherwise. CFR, case fatality rate (%); duration of hospitalisation is presented as median with IQR, interquartile range; h, height; w, weight; congenital abnormalities including Down syndrome and congenital heart disease; SES, socio-economic status; BPH, Biliran Provincial Hospital; EVRMC, Eastern Visayas Regional Medical Center; ONP, Ospital Ng Palawan. (\*)Analysis from February 2014 to March 2016. Past experience of child loss included all causes of death.