

SUPPLEMENTARY TABLE S2. JACKKNIFE SENSITIVITY ANALYSIS OR ROI META-ANALYSIS

Study	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19
FA: TBI < controls																			
Posterior internal capsule, right	X	X	X	X							X	X							
Posterior internal capsule, bilateral														X				X	
Posterior internal capsule, left	X	X	X	X							X	X							
Splenium, bilateral				X	X	X													X
Anterior internal capsule, right	X	X	X								X	X							
Posterior thalamic radiations, left	X										X	X							
MD/ADC: TBI > controls														X	X				X
Splenium of CC, bilateral				X	X		X	X							X	X		X	
Genu of CC, bilateral					X		X	X							X	X	X	X	
Body of CC, bilateral							X							X				X	

1, Adam¹⁹; 2, Arfanakis²⁵; 3, Clark²⁸; 4, Grossman³¹; 5, Inglese³⁰; 6, Kraus³³; 7, Kumar³⁴; 8, Lange³⁵; 9, Ling¹⁸; 10, Little³⁷; 11, Mac Donald (2011)³⁹; 12, Mac Donald (2017)²⁰; 13, Maruta⁴⁰; 14, Matsushita⁴¹; 15, Rutgers⁴⁴; 16, Singh⁴⁵; 17, Sorg⁴⁶; 18, Wilde⁴⁷; 19, Zhang⁴⁸.

An "X" indicates that the results remained significant when the study was removed from analysis. A shaded black box indicates that study was not included in primary analysis.

ROI, region of interest; FA, fractional anisotropy; TBI, traumatic brain injury; MD, mean diffusivity; ADC, apparent diffusion coefficient; CC, corpus callosum.