

**Table S2**

		<b>no-ADT</b>	<b>ADT</b>
<b>age [years]</b>		<b>70±7</b>	<b>71±6</b>
<b>PSA</b> [ng/ml]	<b>initial</b>	<b>23±24</b> (excl. one outlier>200)	<b>21±25</b> (excl. one outlier>200)
	<b>nadir*</b>	<b>0.02 (0.00; 0.13)</b>	<b>0.12 (0.05; 0.30)</b>
<b>Gleason Score</b>	<b>total</b>	<b>7.6±1.0</b>	<b>7.6±1.0</b>
	GS 1	3.6±0.6	3.8±0.6
	GS 2	4.0±0.7	3.8±0.7
<b>Volume [ccm]</b>	<b>lesion</b>	<b>1.8 (0.7; 7.9)</b>	<b>1.3 (0.4; 4.8)</b>
	<b>PZ</b>	<b>7 (7; 17)</b>	<b>9 (6; 13)</b>
	<b>CG</b>	<b>23 (18; 40)</b>	<b>16 (13; 23)</b>

**Table S2** Characteristic parameters of the two patient sub-groups. With exception of the PSA nadir, none of the parameters differed significantly in between patient subgroups.

Mean±SD is given for normal distributed parameters, median (1<sup>st</sup> quartile; 3<sup>rd</sup> quartile) for non-normal distributed ones. \* indicates significant difference ( $p<0.05$ )