

Supplementary Table 1. Frequency of sexual intercourse

	Low-risk group (n=204)					High-risk group (n=133)				
	1–2 times /1 mo	1–2 times /3 mo	1–2 times /6 mo	1–2 times /1 y	None	1–2 times /1 mo	1–2 times /3 mo	1–2 times /6 mo	1–2 times /1 y	None
Age (y)										
60–64	10 (4.9)	1 (0.5)	0 (0.0)	0 (0.0)	4 (2.0)	1 (0.8)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
65–70	23 (11.3)	6 (2.9)	3 (1.5)	5 (2.5)	19 (9.3)	7 (5.3)	2 (1.5)	0 (0.0)	2 (1.5)	10 (7.5)
71–80	27 (13.2)	6 (2.9)	2 (1.0)	2 (1.0)	68 (33.3)	21 (15.8)	5 (3.8)	4 (3.0)	5 (3.8)	45 (33.8)
81–90	1 (0.5)	0 (0.0)	0 (0.0)	0 (0.0)	23 (11.3)	6 (4.5)	1 (0.8)	0 (0.0)	0 (0.0)	21 (15.8)
≥91	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	4 (2.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	3 (2.3)
Sex										
Male	35 (17.2)	11 (5.4)	2 (1.0)	5 (2.5)	44 (21.6)	34 (25.6)	8 (6.0)	4 (3.0)	6 (4.5)	65 (48.9)
Female	26 (12.7)	2 (1.0)	3 (1.5)	2 (1.0)	74 (36.3)	1 (0.8)	0 (0.0)	0 (0.0)	1 (0.8)	14 (10.5)

Values are presented as number (%).