

Supplementary Table 4. Condom use during sexual intercourse

	Low-risk group (n=85)				High-risk group (n=54)			
	Almost always	More than half times	Less than half times	Never used	Almost always	More than half times	Less than half times	Never used
Age (y)								
60–64	0 (0.0)	0 (0.0)	2 (2.4)	10 (11.8)	0 (0.0)	0 (0.0)	0 (0.0)	1 (1.9)
65–70	0 (0.0)	0 (0.0)	0 (0.0)	35 (41.2)	1 (1.9)	0 (0.0)	1 (1.9)	9 (16.7)
71–80	1 (1.2)	0 (0.0)	0 (0.0)	33 (38.8)	2 (3.7)	0 (0.0)	1 (1.9)	32 (59.3)
81–90	0 (0.0)	0 (0.0)	0 (0.0)	1 (1.2)	2 (3.7)	0 (0.0)	0 (0.0)	5 (9.3)
≥91	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Sex								
Male	1 (1.2)	0 (0.0)	1 (1.2)	50 (58.8)	5 (9.3)	0 (0.0)	2 (3.7)	45 (83.3)
Female	0 (0.0)	0 (0.0)	0 (0.0)	29 (34.1)	0 (0.0)	0 (0.0)	0 (0.0)	2 (3.7)

Values are presented as number (%).