

Supplementary Table 13. Preferred institution to receive sexual health education for low-risk group and high-risk group

	Low-risk group (n=204)						High-risk group (n=133)						
	District office	Welfare center	Counseling center	Religious institution	Public health center	Etc.	District office	Welfare center	Counseling center	Religious institution	Public health center	Professional organization	Etc.
Age (y)													
60–64	1 (0.5)	10 (4.9)	0 (0.0)	0 (0.0)	3 (1.5)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.8)	0 (0.0)	0 (0.0)
65–70	5 (2.5)	27 (13.2)	3 (1.5)	0 (0.0)	12 (5.9)	4 (2.0)	3 (2.3)	2 (1.5)	1 (0.8)	1 (0.8)	15 (11.3)	0 (0.0)	0 (0.0)
71–80	0 (0.0)	38 (18.6)	4 (2.0)	0 (0.0)	20 (9.8)	8 (3.9)	6 (4.5)	15 (11.3)	2 (1.5)	1 (0.8)	36 (27.1)	3 (2.3)	7 (5.3)
81–90	0 (0.0)	4 (2.0)	0 (0.0)	0 (0.0)	2 (1.0)	2 (1.0)	0 (0.0)	3 (2.3)	1 (0.8)	0 (0.0)	16 (12.0)	0 (0.0)	1 (0.8)
≥91	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.5)	0 (0.0)	1 (0.8)	1 (0.8)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Sex													
Male	2 (1.0)	41 (20.1)	4 (2.0)	0 (0.0)	19 (9.3)	11 (5.4)	7 (5.3)	19 (14.3)	5 (3.8)	1 (0.8)	56 (42.1)	3 (2.3)	8 (6.0)
Female	4 (2.0)	38 (18.6)	3 (1.5)	0 (0.0)	18 (8.8)	2 (1.0)	2 (1.5)	2 (1.5)	0 (0.0)	0 (0.0)	12 (9.0)	0 (0.0)	0 (0.0)

Values are presented as number (%).