

Supplemental Table 1. Standardized incidence ratios (SIRs) and 95% confidence intervals (CIs) for overall second primary cancer (SPC) risk according to anatomical site	
Anatomical site	SIR (95% CI)
Gum and Other Mouth	0.62 (0.38, 0.96)
Tonsil	0.43 (0.14, 0.99)
Esophagus	0.79 (0.56, 1.08)
Stomach	0.82 (0.67, 0.99)
Small Intestine	1.36 (1.02, 1.79)
Colorectal	1.27 (1.20, 1.35)
Pancreas	0.87 (0.76, 0.98)
Lung and Bronchus	0.81 (0.76, 0.86)
Soft Tissue including Heart	1.44 (1.10, 1.84)
Breast	1.05 (1.01, 1.09)
Vagina	6.12 (4.97, 7.47)
Vulva	1.16 (0.89, 1.49)
Urinary System	1.31 (1.21, 1.42)
Urinary Bladder	1.20 (1.07, 1.35)
Kidney	1.46 (1.29, 1.64)
Endocrine System	1.44 (1.27, 1.63)
Lymphocytic Leukemia	0.73 (0.57, 0.91)
Acute Non-Lymphocytic Leukemia (ANLL)	1.34 (1.09, 1.62)
Myeloid and Monocytic Leukemia	1.23 (1.02, 1.47)
Acute Myeloid Leukemia	1.37 (1.10, 1.68)