

Supplementary Table 1. Comparing various clinical characteristics of survivors and non-survivors among RRT patients (n=65)

Characteristics, n (%)	Survivors n=19 (29.2%)	Non-survivors n=46 (70.8%)	<i>p value</i>
Age, mean ± SD	61.9 ± 17.3	58.9± 15.7	0.48
Men	11 (57.9)	34 (73.9)	0.20
CCI, median (IQR)	4 (2-5)	4 (2-6)	0.70
Pre-arrest comorbidities			
Myocardial Infarction	2 (10.5)	2 (4.4)	0.57
Heart Failure	4 (21.1)	16 (34.8)	0.28
Hypertension	13 (68.4)	28 (60.9)	0.57
Renal Disease	7 (36.8)	17 (37.0)	0.99
NIDDM	6 (31.6)	8 (17.4)	0.32
Liver Disease	0 (0)	5 (10.9)	0.31
Cerebrovascular Disease	1 (5.3)	3 (6.5)	1.00
PVD	2 (10.5)	6 (13.0)	1.00
Obesity (BMI>30 m/kg ²)	2 (10.5)	6 (13.0)	1.00
Dementia	1 (5.3)	2 (4.4)	1.00
Smoking status			0.20
Non-smoker	11 (57.9)	24 (57.1)	
Former-smoker	7 (36.8)	9 (21.4)	
Current-smoker	1 (5.3)	9 (21.4)	
Pre-arrest CPC			0.60
- CPC 1	14 (73.7)	29 (63.0)	

- CPC 2	4 (21.1)	11 (23.9)	
- CPC 3	1 (5.3)	6 (13.0)	
in-hospital arrests	12 (63.2)	23 (50.0)	0.33
Witnessed	16 (84.2)	37 (80.4)	1.00
CPR			0.09
-No CPR	0 (0)	10 (21.7)	
-Yes, by a bystander	3 (15.8)	6 (13.0)	
-Yes, with medical personnel	16 (84.2)	30 (65.2)	
Defibrillation use	7 (38.9)	17 (37.8)	0.94
Initial Rhythm			0.69
- VT/VF/AED Advised Shock	8 (42.1)	12 (30.8)	
- PEA	7 (36.8)	18 (46.2)	
- Asystole	4 (21.1)	9 (23.1)	
ROSC, mean ± SD	16.9 ± 15.7	27.5 ± 29.5	0.21
Temperature Management	13 (68.4)	40 (87.0)	0.16
Dialysis requirements at discharge (for survivors only)	7 (36.8)	--	--

SD: standard deviation

BMI: Body Mass Index

IQR: Interquartile Range

NIDDM: Non-insulin-dependent diabetes mellitus

PVD: Peripheral vascular disease

CPC: Cerebral performance score

CPR: Cardiopulmonary resuscitation

VT/VF: Ventricular tachycardia / ventricular fibrillation

AED: Automated external defibrillators

PEA: Pulseless electrical activity

ROSC: Return of Spontaneous Circulation