

Table S7: Moderate/heavy infection prevalence, combined nutrition plus WSH vs. WSH and nutrition interventions

Arm	N	Prevalence	Prevalence ratio			Prevalence difference		
			Unadjusted	Adjusted <sup>a</sup>	IPCW <sup>b</sup>	Unadjusted	Adjusted <sup>a</sup>	IPCW <sup>b</sup>
<b>Ascaris</b>								
Nutrition + WSH	933	6.1%						
WSH	941	3.3%	1.85 (1.09, 3.16)	2.02 (1.21, 3.39)	2.02 (1.18, 3.47)	2.81 (0.40, 5.23)	3.15 (0.90, 5.41)	3.16 (0.77, 5.55)
Nutrition	863	4.2%	1.46 (0.89, 2.42)	1.42 (0.88, 2.29)	1.46 (0.87, 2.45)	1.94 (-0.67, 4.55)	1.79 (-0.69, 4.26)	1.96 (-0.76, 4.68)
<b>Hookworm</b>								
Nutrition + WSH	933	0.0%						
WSH	941	0.2%	- <sup>c</sup>	- <sup>c</sup>	- <sup>c</sup>	-0.21 (-0.50, 0.08)	-0.26 (-0.52, 0.00)	-0.22 (-0.54, 0.09)
Nutrition	863	0.3%	- <sup>c</sup>	- <sup>c</sup>	- <sup>c</sup>	-0.35 (-0.74, 0.05)	-0.36 (-0.74, 0.02)	-0.35 (-0.72, 0.03)
<b>Trichuris</b>								
Nutrition + WSH	933	1.1%						
WSH	941	0.7%	1.44 (0.50, 4.17)	1.39 (0.50, 3.86)	1.41 (0.54, 3.68)	0.33 (-0.59, 1.25)	0.30 (-0.58, 1.18)	0.31 (-0.53, 1.15)
Nutrition	863	0.5%	2.31 (0.95, 5.65)	2.26 (0.94, 5.44)	2.27 (0.90, 5.72)	0.61 (-0.08, 1.29)	0.59 (-0.07, 1.25)	0.60 (-0.13, 1.33)
<b>Any STH</b>								
Nutrition + WSH	933	6.6%						
WSH	941	3.8%	1.74 (1.06, 2.84)	1.88 (1.17, 3.01)	1.87 (1.14, 3.07)	2.82 (0.30, 5.34)	3.17 (0.84, 5.51)	3.15 (0.66, 5.65)
Nutrition	863	4.8%	1.40 (0.85, 2.29)	1.33 (0.83, 2.11)	1.39 (0.85, 2.29)	1.89 (-0.98, 4.77)	1.60 (-1.09, 4.29)	1.89 (-1.01, 4.80)

<sup>a</sup> Adjustment covariates considered include ID of the lab staff member who performed the Kato-Katz analysis, month of measurement, child age, sex and birth order since this information is not available for individuals lost to follow-up. An indicator variable distinguishing index vs. non-index child status was included as a proxy for age.

<sup>b</sup> Inverse probability of censoring weighting. Adjustment covariates considered include the variables above except for ID of the lab staff member who performed the Kato-Katz analysis, month of measurement, child age, sex and birth order since this information is not available for individuals lost to follow-up. An indicator variable distinguishing index vs. non-index child status was included as a proxy for age.

<sup>c</sup> Could not calculate due to sparse data.