

Article

Solid Matrix-Supported Supercritical CO₂ Enhances Extraction of γ -Linolenic Acid from the Cyanobacterium *Arthrospira (Spirulina) platensis* and Bioactivity Evaluation of the Molecule in Zebrafish

Xiaohong Yang ^{1,5}, Yi Li ^{2,6}, Yanhua Li ¹, Ding Ye ^{2,6}, Li Yuan ¹, Yonghua Sun ^{2,6}, Danxiang Han ^{1,4,*} and Qiang Hu ^{1,2,3,4,7,*}

¹ Center for Microalgal Biotechnology and Biofuels, Institute of Hydrobiology, Chinese Academy of Sciences, Wuhan 430072, China; xhyang@ihb.ac.cn (X.Y.); yanhuali@ihb.ac.cn (Y.L.); liyuan@ihb.ac.cn (L.Y.)

² State Key Laboratory of Freshwater Ecology and Biotechnology, Institute of Hydrobiology, Chinese Academy of Sciences, Wuhan 430072, China; liyi_0831@126.com (Y.L.); yeding@ihb.ac.cn (D.Y.); yhsun@ihb.ac.cn (Y.S.)

³ Laboratory for Marine Biology and Biotechnology, Qingdao National Laboratory for Marine Science and Technology, Qingdao 266237, China

⁴ Key Laboratory for Algal Biology, Institute of Hydrobiology, Chinese Academy of Sciences, Wuhan 430072, China

⁵ College of Life Sciences, University of Chinese Academy of Sciences, Beijing 100049, China

⁶ College of Advanced Agricultural Sciences, University of Chinese Academy of Sciences, The Innovative Academy of Seed Design, Chinese Academy of Sciences, Beijing 100049, China

⁷ Beijing Key Laboratory of Algae Biomass, Microalgae Biotechnology Center, SDIC Biotech Investment Co., LTD., State Development & Investment Corp., Beijing 100142, China

* Correspondence: huqiang@ihb.ac.cn (Q.H.); danxianghan@ihb.ac.cn (D.H.)




Figure S1. Comparison of the states of mixing porous materials with algal powders and the microscopic images of residues (Scale bar 50 μ m).




Figure S2. The chromatograms of fatty acid profiles obtained from the solid matrix-supported supercritical CO₂ (SMSSC-CO₂), ethanol, Bligh and Dyer, and in situ transesterification methods.