

Supplementary Table. Proportion of PLNM among patients who received prostatectomy and who didn't receive prostatectomy. Values represent n (%) per year.

<b>Year:</b>	<b>2004</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>	<b>2013</b>	<b>2014</b>
<b>All men 50 years and older</b>											
RP	307(2.0)	307(2.1)	332(2.0)	397(2.1)	401(2.2)	417(2.2)	457(2.5)	467(2.5)	508(3.4)	549(3.9)	613(4.7)
Non-RP	215(0.7)	220(0.7)	215(0.6)	236(0.7)	239(0.7)	251(0.8)	294(0.9)	284(0.9)	255(0.9)	296(1.0)	372(1.4)
<b>Age Stratified</b>											
<b>50 to 74 years old</b>											
RP	299(2.0)	290(2.1)	316(1.9)	376(2.1)	379(2.1)	398(2.2)	437(2.4)	437(2.4)	481(3.3)	518(3.8)	583(4.7)
Non-RP	176(0.8)	190(0.9)	179(0.8)	185(0.8)	183(0.8)	195(0.8)	226(0.9)	209(0.8)	187(0.9)	214(1.0)	270(1.4)
<b>75 years and older</b>											
RP	8(2.0)	17(4.4)	16(3.7)	21(4.2)	22(4.9)	19(4.2)	20(3.9)	30(5.6)	27(6.1)	31(7.0)	30(7.0)
Non-RP	39(0.4)	30(0.3)	36(0.3)	51(0.5)	56(0.6)	56(0.6)	68(0.8)	75(0.9)	68(1.0)	82(1.3)	102(1.7)

P trend<0.05 for all groups from Cochran-Armitage trend test