

S2 Table. Samples weight (grams) of each group destined for qPCR.

	Parotid			Submandibular			Sublingual		
	Saline	Losartan	Isoproterenol	Saline	Losartan	Isoproterenol	Saline	Losartan	Isoproterenol
	217.5	221.4	394.7	153.8	169.2	232.7	38.4	39.1	34.3
	327.2	225.8	394.6	176.4	175.5	272.4	26.7	27.7	33.5
	235.5	185.1	475.3	159.7	162.1	342.2	35.8	27.7	29.8
	206.4	233	625.1	179.1	179.1	328.2	36.7	35.8	34.2
	265.6	175.1	551.5	147.8	161.0	325.9	35.7	39.3	43.8
	234.7	310.9	*	163.1	200.4	*	29	27.6	*
	267.5	216.1	*	183.1	183.7	*	44	34	*
Mean	250.6^a	223.9^a	488.2^b	166.1^a	175.8^a	300.2^b	35.1	33.0	35.1
SD	40.6	43.9	100.5	13.5	13.7	46.1	5.7	5,30	5,10

Data are means \pm standard deviation. One-way ANOVA/Tukey's multiple comparison tests were performed. Differences were considered statistically significant when $p < 0.05$. Different letters (a or b) indicate a statistically significant difference between groups, whereas equal letters indicate the absence of such differences. Letters were not used when differences were not significant between saline, losartan and isoproterenol. It is important to mention that the ipsilateral gland was sent for immunohistochemistry and is not included in the measurements of this table.