

Supplementary Table S8
Sources of genomic DNA used for HCAR amplification

species		source
<i>Gorilla gorilla</i>	Western Gorilla	Dr T. Haaf, MPI Molecular Genetics Berlin, Germany; Dr L. Vigilant, MPI, Leipzig, Germany
<i>Homo sapiens</i>	human	Dr M. Stoneking, MPI, Leipzig, Germany; D. Sere, MPI, Leipzig, Germany
<i>Mus musculus</i>	house mouse	Dr A. Orth, University of Montpellier, France
<i>Nomascus leucogenys</i>	white-cheeked Gibbon	Dipl. Biol. C. Roos, Primate Center Göttingen, Germany
<i>Pan paniscus</i>	bonobo	Dr W. Enard, MPI, Leipzig, Germany; Dr L. Vigilant, MPI, Leipzig, Germany
<i>Pan troglodytes</i>	chimpanzee	Dr T. Haaf, MPI Molecular Genetics Berlin, Germany; Dr W. Enard, MPI, Leipzig, Germany
<i>Pongo pygmaeus</i>	Orangutan	Dr M. Rocchi, University of Bari, Italy; Dr W. Enard, MPI, Leipzig, Germany
<i>Symphalangus syndactylus</i>	siamang	Dipl. Biol. C. Roos, Primate Center Göttingen, Germany; Dr T. Haaf, MPI Molecular Genetics Berlin, Germany